

## AMELIORATION BY MELATONIN OF CHROMOSOMAL ANOMALIES INDUCED BY ARSENIC AND/OR FLUORIDE IN HUMAN BLOOD LYMPHOCYTE CULTURES<sup>a</sup>

Mandava V Rao,<sup>b</sup> Hemlata Tiwari

Ahmedabad, India

**SUMMARY:** Standard cytochemical methods were used to investigate the ameliorative effect of melatonin (0.2 mM) on chromosomal aberrations in human lymphocyte cultures induced by arsenic (As<sub>2</sub>O<sub>3</sub>, 1.4 μM) and/or fluoride (NaF, 34 μM). As<sub>2</sub>O<sub>3</sub> and/or NaF generated a significant increase in the incidence of chromosomal aberrations as compared to control levels. Combined treatment with As<sub>2</sub>O<sub>3</sub> and NaF induced more chromosomal aberrations and aneuploidy than either reagent individually. Melatonin supplements brought about a significant decrease in the number of aberrations, with the percentage of amelioration varying between 53% and 88%. This reduction by melatonin of genotoxic effects exerted by As and/or F is probably attributable to its protective antioxidant action.

Keywords: Arsenic and chromosomes; Chromosomal aberrations; Lymphocyte cultures; Melatonin and chromosomes; Fluoride and chromosomes; Oxidative stress.

### INTRODUCTION

Arsenic and fluoride are well-known water contaminants, and their toxicity on humans has been widely studied. Arsenic is a known human carcinogen and induces cancers of lung, skin, bladder, kidneys and liver.<sup>1</sup> It is introduced into water through dissolution of rocks, minerals, and ores, from industrial effluents including mining wastes, and from atmospheric deposition.<sup>2</sup> Inorganic As<sup>+3</sup> does not appear to be a mutagen in standard assays,<sup>3</sup> whereas As<sup>+3</sup> exposure *in vitro* produces chromosomal aberrations, DNA protein cross links, and SCE (sister chromatid exchanges) in hamster embryo cells<sup>4,5</sup> and in human lymphocytes.<sup>6</sup>

On the other hand, fluoride (F as F<sup>-</sup>) is also known to be a major environmental contaminant. In India more than 20 states are affected by F toxicity, Gujarat being the most heavily impacted state.<sup>7</sup> With respect to genotoxic effects of F, there has been much confusion and a lack of adequate information. Joseph et al.<sup>8</sup> reported greater rates of SCE and chromosomal aberrations in people living in F-endemic villages than in nonendemic villages. Some authors, however, report that F does not induce genotoxic effects.<sup>9</sup> Earlier studies at our institution have reported mitigation of As<sup>+3</sup> and/or F genotoxicity by vitamins as antioxidants in the Indian population.<sup>10</sup>

Melatonin (N-acetyl-5-methoxytryptamine), the major secretory product of the pineal gland, is involved in the regulation of circadian rhythm and seasonal changes in vertebrate physiology.<sup>11</sup> It is identified as a powerful direct free radical scavenger<sup>12</sup> as well as an indirect antioxidant.<sup>13</sup> It has also been shown to reduce molecular damage effectively under conditions of elevated oxidative stress.<sup>14</sup> Melatonin, in fact, was found to exhibit better antioxidative properties under *in vitro* conditions than similar concentrations of α-tocopherol.<sup>15</sup> In view of its strong antioxidant activity, an investigation has now been undertaken of melatonin

<sup>a</sup>This paper is dedicated to our beloved teacher, the late Prof NJ Chinoy (d. 8 May 2006).

<sup>b</sup>For correspondence: Dr MV Rao, Prof and Head, Department of Zoology, University School of Sciences, Gujarat University, Ahmedabad 380 009, India; Email: zooldeptgu@satyam.net.in

for its ability to mitigate genotoxic effects of As and F in human blood cultures obtained from volunteers in India.

### MATERIALS AND METHODS

*Subjects:* Venous blood was collected in sterile heparinised syringes from healthy, consenting 20–25 year old non-smoking residents of Ahmedabad, India.

*Peripheral blood lymphocyte culture (PBLC):* PBLC was then performed by the method of Perry and Wolff.<sup>16</sup> To 7 mL of RPMI-1640 (HiMedia, Mumbai, pH 7.4) previously supplemented with 7% fetal calf serum (FCS), antibiotics (benzyl penicillin and streptomycin) and 100  $\mu$ L of phytohemagglutinin [PHA], 5mg/5mL distilled water; Sigma-Aldrich, USA), nine drops of blood were added. After 48 hr of incubation, As<sub>2</sub>O<sub>3</sub> (HiMedia, Mumbai) and NaF (Qualigens Fine Chemicals, Mumbai), with or without melatonin (HiMedia, Mumbai), were added. Harvesting was done after 24 hr.

Cultures were divided into eight groups. Group I was the control i.e., blood culture without any treatment. In Group II, lymphocytes were treated with As (As<sub>2</sub>O<sub>3</sub>; 1.4  $\mu$ M), Group III with F (NaF; 34  $\mu$ M), and Group IV with a combination of the same concentrations of As<sub>2</sub>O<sub>3</sub> and NaF. Antioxidant Groups were melatonin (0.2 mM) supplemented with As (Group V), F (Group VI) and As+F (Group VII). Group VIII contained only melatonin added to the blood culture.

After 69 hr, 15  $\mu$ L of colchicine (1 mg/5 mL distilled water, HiMedia, Mumbai) was added for 30 min to arrest cells at the metaphase stage. The pellet obtained after centrifugation was treated with hypotonic solution (0.075 M KCl) for 20 min at 37°C. These cells were then fixed by chilled 1:3 acetic acid:methanol fixative. Slides were prepared from the cell suspension obtained after two washes with fixative. These slides were then stained with 2% Geimsa stain and observed under the microscope at 100X for scoring structural and numerical aberrations.

*Analysis of parameters:* One hundred plates per group were analysed under the microscope for chromosome and chromatid breaks and gaps and number of chromosomes. Plates with a chromosome number greater than 44 were used for scoring structural and numerical aberrations. Percentage of amelioration was calculated by using the following formula:

$$\frac{(\text{Pro-oxidant Group} - \text{Respective Antioxidant Group})}{(\text{Pro-oxidant Group} - \text{Control})} \times 100$$

*Statistical analysis:* Results are expressed as mean  $\pm$  SE. All the treated groups were compared with the control group, and the melatonin-supplemented groups were compared with their respective pro-oxidant groups by Student's t-test. P values less than 0.05 were considered to be significant.

### RESULTS

Addition of As (as As<sub>2</sub>O<sub>3</sub>) and/or F (as NaF) to the peripheral blood cultures showed a highly significant ( $p < 0.001$ ) increase in the number of total chromosomal aberrations as compared to the control cultures. On the other hand,

co-culturing of melatonin with As and/or F demonstrated a highly significant ( $p < 0.001$ ) decline in the number of induced chromosomal aberrations in comparison to the respective pro-oxidant group. Melatonin exhibited 52% and 74% amelioration from individual toxicity induced by As and F, respectively, and 64% amelioration from combined toxicity, for chromosomal aberrations (Table 1).

As seen in Table 2, the presence of As alone and in combination with F led to a very significant increase in the number of hypoploids ( $p < 0.001$ ) as compared to the control. F alone also caused a marked increase in the number of hypoploids ( $p < 0.05$ ). Addition of melatonin to the As and F treated cultures individually as well as to their combination produced significant reductions in the number of hypoploids with values that are comparable to those of the control culture. The amelioration was 79% and 88% from the toxicity of As and F, respectively ( $p < 0.05$  and  $p < 0.001$ ), and 84% from their combination ( $p < 0.001$ ). Melatonin alone had essentially no effect and gave readings comparable to the control values (Tables 1 and 2).

**Table 1.** Effect of melatonin (M) on As- and/or F-induced chromosome and chromatid breaks and gaps per 100 plates in human lymphocytes after 24-hr exposure for 5 individuals

Group	Chromatids		Chromosomes		Mean <sup>a</sup> ± SE	Values in percent	t-test with control <sup>b</sup>	t-test with resp. Groups <sup>c</sup>	Amelioration (%)
	Breaks	Gaps	Breaks	Gaps					
Control	1.20	1.20	0.80	0.80	4.00 ± 0.80	100			
As	30.60	10.80	9.00	5.80	56.20 ± 1.57	1405	29.73 <sup>†</sup>		
F	13.60	6.60	5.60	5.80	31.60 ± 0.96	790	22.21 <sup>†</sup>		
As+F	34.80	15.60	7.60	6.80	64.80 ± 1.82	1620	25.31 <sup>†</sup>		
As+M	15.20	5.60	4.80	3.20	28.80 ± 1.66	720	12.96 <sup>†</sup>	12.01 <sup>†</sup>	53
F+M	5.40	3.00	1.80	1.00	11.20 ± 1.07	280	3.41 <sup>†</sup>	14.20 <sup>†</sup>	74
As+F+M	14.60	3.60	4.60	3.20	26.00 ± 1.43	650	10.03 <sup>†</sup>	16.73 <sup>†</sup>	64
M	3.00	1.4	0.80	0.8	6.00 ± 0.56	150	1.65 <sup>ns</sup>		

<sup>a</sup>Mean indicates total structural aberrations; <sup>b</sup>all groups were compared with control; <sup>c</sup>antioxidants groups were compared with respective pro-oxidant (As and/or F) group.  
<sup>†</sup> $p < 0.001$ ; <sup>†</sup> $p < 0.01$ ; <sup>ns</sup>not significant.

**Table 2.** Effect of melatonin on As and/or F induced numerical aberrations (hypoploidy) in human lymphocyte chromosomes after 24-hr exposure for 5 individuals

Groups	Mean <sup>a</sup> ± SE	Values in percent	t-test with control <sup>b</sup>	t-test with resp. groups <sup>c</sup>	Amelioration (%)
Control	11 ± 1.00	100			
As	23 ± 1.21	209	7.39*		
F	18 ± 1.12	164	4.27 <sup>†</sup>		
As+F	26 ± 1.05	236	10.45*		
As+M	14 ± 1.05	127	1.65 <sup>ns</sup>	6.79 <sup>†</sup>	79
F+M	12 ± 0.89	109	0.60 <sup>ns</sup>	4.82 <sup>†</sup>	88
As+F+M	14 ± 1.00	127	1.70 <sup>ns</sup>	11.54 <sup>†</sup>	84
M	10 ± 1.03	90.9	-0.70 <sup>ns</sup>		

<sup>a</sup>Mean indicates number of numerical aberrations; <sup>b</sup>all groups were compared with control; <sup>c</sup>antioxidants groups were compared with respective pro-oxidant (As and/or F) group.

\* $p < 0.001$ ; <sup>†</sup> $p < 0.05$ ; <sup>ns</sup>not significant.

## DISCUSSION

Results of the present study revealed an increase in the number of structural as well as numerical chromosome and chromatid aberrations induced by As and/or F, indicating their genotoxicity. This induced genotoxicity is probably mediated by induction of oxidative stress and depletion of glutathione<sup>17-19</sup> as a result of the action of these two pro-oxidants. Electron spin resonance (ESR) analysis using

spin trapping with 5,5-dimethyl-1-pyrroline-N-oxide also suggests that reactive oxygen species (ROS) participate in genotoxicity of As.<sup>20</sup> Thus, Oya et al.<sup>21</sup> found induction by As of chromosomal aberrations by hydrogen peroxide. Similarly, Eguchi et al.<sup>22</sup> observed increased frequency of chromosomal aberrations by As as observed here. Their study also revealed an increased frequency of aneuploidy on addition of As, in agreement with findings on HFW cells by Yih et al.<sup>23</sup> It has been documented that loss of p53 activity leads to aneuploidy along with gene amplification, and since As is known to alter p53 activity,<sup>24</sup> this might be the possible cause of the observed hypoploidy.

It is known that F<sup>-</sup> affects enzymatic activities, and this effect could delay mitotic cycles causing chromosomal breakages.<sup>25</sup> Similarly, reports are available that F increases chromosomal damage,<sup>8,26,27</sup> corroborating our data. Ardema et al.<sup>28</sup> proposed that NaF-induced aberrations might occur by an indirect mechanism involving inhibition of DNA synthesis/repair due to the formation of hydrogen bonds with nitrogen bases. Further, F also affects polymerase activity involved in DNA replication, since F has great affinity to Ca<sup>+2</sup>, Mg<sup>+2</sup>, and phosphate ions.<sup>10</sup> Chinoy et al.<sup>29</sup> also found increased aneuploidy in lymphocyte cultures after addition of NaF, again confirming the interaction of F with DNA nucleotides. In the present study, genotoxic effects of the combined treatment of As and F were more pronounced than with individual treatments, as might be expected. Likewise, genotoxic effects of As and/or F on human blood cultures of the Indian population have been reported by Nair et al.<sup>10</sup>

Melatonin has been shown to protect membrane lipids, nuclear DNA, and protein from oxidative damage induced by variety of free radical generating agents.<sup>30,31,32</sup> Vijayalaxmi et al.<sup>33</sup> reported that addition of melatonin reduced the incidence of primary DNA damage and chromosomal aberrations as well as the induction of micronuclei due to increased oxidative stress in irradiated human blood lymphocyte cultures. Melatonin, has also been used against oxidative stress-mediated DNA damage by chromium, cyclophosphamide, and hydrogen peroxide.<sup>34,35</sup> Since melatonin is subcellularly widespread, it might allow maximum interaction with all molecules thus protecting against ROS in both aqueous and lipid environment.<sup>13</sup> The present study also indicated a decrease in chromosomal aberrations on addition of melatonin to cell cultures exposed to pro-oxidant As and F. This ameliorative effect of melatonin could be attributed to scavenging hydroxyl radical, stimulating antioxidative enzymes, inhibiting pro-oxidative enzymes,<sup>36,37</sup> possibly by involvement in the formation of As<sup>+3</sup>/F<sup>-</sup> complexes, thereby interfering with normal DNA and protein interactions, leading to chromosomal aberrations.

This study thus demonstrates that melatonin supplementation provides protection against chromosomal anomalies induced by As and/or F under *in vitro* condition

#### ACKNOWLEDGEMENT

This work was supported by University Grants Commission (UGC), New Delhi, in the form of a Major Research Project to MVR and Junior Research Fellowship (JRF) to HT.

## REFERENCES

- 1 Tom KH, Metka F. Role of oxidative damage in genotoxicity of arsenic. *Free Radic Biol Med* 2004;37(5):574-81.
- 2 ATSDR. US Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease research. Toxicological profile for arsenic (update). 2000.
- 3 Hughes MF. Arsenic toxicity and potential mechanism of action. *Toxicol Lett* 2002;133:1-16.
- 4 Lee TC, Oshimura N, Barrett JC. Comparison of arsenic induced cell transformation, cytotoxicity, mutation and cytogenetic effects in Syrian hamster embryo cells in culture. *Carcinogenesis* 1985;6(10):1421-6.
- 5 Dong JT, Luo XM. Arsenic induced breaks associated with DNA-protein cross links in human fetal fibroblast. *Mutat Res* 1993;302:97-102.
- 6 Jha AN, Noditi N, Nilsson R, Natrajan AT. Genotoxic effects of sodium arsenite on human cells. *Cancer Lett* 1992;21:141-7.
- 7 Susheela AK. Fluorosis management programme in India. *Curr Sci* 1999;77:1250-6.
- 8 Joseph H, Ghadia PK. Sister chromatid exchange frequency and chromosome aberrations in residents of fluoride endemic regions of South Gujarat. *Fluoride* 2000;33:154-58.
- 9 Tong CC, McQueen CA, Brat SV, Williams GM. The lack of genotoxicity of sodium fluoride in a battery of cellular tests. *Cell Biol Toxicol* 1988;4(2):173-86.
- 10 Nair SB, Jhala DD, Chinoy NJ. Mitigation of genotoxic effects of fluoride and arsenic by ascorbic acid in human lymphocyte cultures. *Fluoride* 2004;37(4):249-56.
- 11 Antonio C, Juan RC, Pedro A, Patricia JL, Sofia GM, Russel JR, et al. Evidence of melatonin synthesis by human lymphocyte and its physiological significance: possible role as intracrine, autocrine and/or paracrine substance. *FASEB J* 2004;18:537-9.
- 12 Tan DX, Reiter RJ, Manchester LC, Yen MT, El-Sawi M, Sainz RM, et al. Chemical and physical properties and potential mechanisms; Melatonin as a broad spectrum antioxidant and free radical scavenger. *Curr Top Med Chem* 2002;2:181-98.
- 13 Karbownik M, Reiter RJ. Antioxidative effects of melatonin in protection against cellular damage caused by ionizing radiation. *Proc Soc Exp Biol Med* 2000;22:225-9.
- 14 Reiter RJ. Oxidative damage in the central nervous system: Protection by melatonin. *Prog Neurobiol* 1998;56:359-84.
- 15 Ilhami G, Buyukokuroglu ME, Oktay M, Kufrevioglu OI. On the *in vitro* antioxidative properties of melatonin. *J Pineal Res* 2002;33:167-71.
- 16 Perry P, Wolff S. New Geimsa method for the differential staining of sister chromatids. *Nature* 1974;251:156-8.
- 17 Lee TC, Ko JL, Jau KY. Differential cytotoxicity of sodium arsenite in human fibroblast and Chinese hamster ovary cells. *Toxicology* 1989;56:289-99.
- 18 Nordenson I, Beckman L. Is the genotoxic effect of arsenic mediated by oxygen free radicals? *Human Hered* 1991;41:71-3.
- 19 Wang TS, Hsu TY, Chung CH, Wang AS, Bau DT, Jau KY. Arsenite induces oxidative DNA adducts and DNA protein cross links in mammalian cells. *Free Radic Biol Med* 2001;31:321-30.
- 20 Nesnow S, Roop BC, Lambert G, Kadiiska M, Mason RP, Cullen WR, et al. DNA damage induced by methylated trivalent arsenicals is mediated by reactive oxygen species. *Chem Res Toxicol* 2002;15:1627-34.
- 21 Oya OY, Kaise T, Ochi T. Induction of chromosomal aberrations in cultured human fibroblasts by inorganic and organic arsenic compounds, the different roles of glutathione in such induction. *Mutat Res* 1996;357:123-9.
- 22 Eguchi N, Kuroda K, Endo G. Metabolites of arsenic induced tetraploids and mitotic arrest in cultured cells. *Arch Environ Contam Toxicol* 1997;32:141-5.
- 23 Yih LH, Ho IC, Lee TC. Sodium arsenite disturbs mitosis and induces chromosome loss in human fibroblast. *Cancer Res* 1997;57:5051-9.
- 24 Rossman TG. Mechanism of arsenic carcinogenesis: an integrated approach. *Mutat Res* 2003;533:37-65.

- 25 Bogin EM, Abrams M, Avidar Y, Israeli B. Effect of fluoride on enzymes from serum, liver, kidney, skeletal and heart muscles of mice. *Fluoride* 1976;9:42-56.
- 26 Meng Z, Zang B. Chromosomal aberration and micronuclei in lymphocyte of workers at phosphate fertilizer factory. *Mutat Res* 1997;393:283-8.
- 27 Tustsui T, Suzuki N, Ohmori M. Sodium fluoride induced morphological and neoplastic transformation, chromosome aberrations, sister chromatid exchanges and unscheduled DNA synthesis in cultured Syrian hamster embryo cells. *Cancer Res* 1984;44:938-41.
- 28 Ardema MJ, Gibson DP, LeBoeuf RA. Sodium fluoride induced chromosome aberrations in different stages of cell cycle: a proposed mechanism. *Mutat Res* 1989;223(2):191-203.
- 29 Chinoy NJ, Jhala DD, Nair SB. Micronuclei and aneuploidy frequencies in human peripheral lymphocytes treated with sodium fluoride [abstract]. *Fluoride* 2002;35(4):251.
- 30 Reiter RJ, Melchiorri D, Sewerynek E, Poeggeler B, Barlow WL, Chuang J, et al. A review of the evidence supporting melatonin's role as an antioxidant. *J Pineal Res* 2001;18:1-11.
- 31 Romero P, Osuna C, Garcia PA, Carrillo VA, Guerrero JM. The pineal secretory product melatonin reduces hydrogen peroxide induced DNA damage in U937 cells. *J Pineal Res* 1999;26:227-35.
- 32 Wakatsuki A, Okatani Y, Ikenova N, Shinohara K, Watanabe K, Fukaya T. Melatonin protects against oxidised low density lipoprotein- induced inhibition of nitric oxide production in human umbilical artery. *J Pineal Res* 2001;31:281-8.
- 33 Vijayalaxmi, Reiter RJ, Tan DX, Herman TS, Thomas CR. Melatonin as a radio protective agent: A review. *Int J Radiat Oncol Biol Phys* 2004;59(3):639-53.
- 34 Salvia RD, Fiore M, Alitti T, Festa F, Ricordy R, Cozzi R. Inhibitory action of melatonin on H<sub>2</sub>O<sub>2</sub> and cyclophosphamide induced DNA damage. *Mutagenesis* 1999;14(1):107-12.
- 35 Wenbo Q, Reiter RT, Tan DX, Garcia JJ, Manchester LC, Karbownik M, Calvo JR. Chromium (III)-induced 8-hydroxy deoxyguanosine in DNA and its reduction by antioxidants: Comparative effects of melatonin, ascorbate and vitamin E. *Environ Health Perspect* 2000;108:399-402.
- 36 Martin M, Macias M, Escames G, Leon J, Acuna-Castroviejo D. Melatonin but not vitamins C and E maintains glutathione homeostasis in t-butyl hydroperoxide-induced mitochondrial oxidative stress. *FASEB J* 2000;14:1677-9.
- 37 Leon J, Acuna-Castroviejo D, Sainz RM, Mayo JC, Tan DX, Reiter RJ. Melatonin and mitochondrial function. *Life Sci* 2004;75(7):765-90.