



C. elegans as a model in developmental neurotoxicology

Joanna A. Ruszkiewicz^{a,*}, Adi Pinkas^a, Mahfuzur R. Miah^a, Rebecca L. Weitz^a,
Michael J.A. Lawes^a, Ayodele J. Akinyemi^{a,b}, Omamuyovwi M. Ijomone^{a,c}, Michael Aschner^a

^a Department of Molecular Pharmacology, Albert Einstein College of Medicine, Bronx, New York, United States

^b Department of Biochemistry, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria

^c Department of Anatomy, School of Health and Health Technology, Federal University of Technology Akure (FUTA), Nigeria

ARTICLE INFO

Keywords:

Neurodevelopment
Neurotoxicity
C. elegans
Manganese
Mercury
Pesticides

ABSTRACT

Due to many advantages *Caenorhabditis elegans* (*C. elegans*) has become a preferred model of choice in many fields, including neurodevelopmental toxicity studies. This review discusses the benefits of using *C. elegans* as an alternative to mammalian systems and gives examples of the uses of the nematode in evaluating the effects of major known neurodevelopmental toxins, including manganese, mercury, lead, fluoride, arsenic and organo-phosphorus pesticides. Reviewed data indicates numerous similarities with mammals in response to these toxins. Thus, *C. elegans* studies have the potential to predict possible effects of developmental neurotoxicants in higher animals, and may be used to identify new molecular pathways behind neurodevelopmental disruptions, as well as new toxicants.

1. Neurodevelopmental toxicity studies – a need for alternative models

The developing brain is exceptionally sensitive to toxic chemicals – the levels which lead to adverse effects are often lower than those effective in adults, and even minimal exposures, generally considered as safe, might be harmful. Moreover, only a few chemicals have been studied and identified as toxic to developing brain, and safe dosages have only been estimated. With over 1000 chemicals currently identified as toxic to animals, and > 200 recognized as neurotoxic to the adult human, only 12 are characterized as toxic to the developing human brain (Grandjean and Landrigan, 2014). This is mostly due to the fact that neurotoxicity is usually discovered during incidents of acute poisoning in adults, specifically from occupational exposure or suicide attempts, which provides clear association between exposure and adverse effects. In contrast, in developmental neurotoxicity the exposure occurs sometimes years before any neurological defects might be identified, and therefore the identification of new hazards is often much slower.

As proposed by Grandjean & Landrigan, the current list of developmental neurotoxins includes arsenic (As), lead (Pb), methylmercury (MeHg), manganese (Mn), fluoride (F), ethanol (EtOH), toluene (PhMe), tetrachloroethylene (PERC), polybrominated diphenyl ethers (PBDEs), organophosphate pesticides (chlorpyrifos) and dichlorodiphenyltrichloroethane (DDT)/dichlorodiphenyldichloro-ethylene (DDE) (Grandjean

and Landrigan, 2014). These compounds have been shown to disrupt neurogenesis, brain cell proliferation, migration, synapse formation and myelination during development (Hu et al., 2014), leading to physiological and behavioral deficits which are often permanent and untreatable. Impaired central nervous system (CNS) function might comprise cognitive, motor, language and affective disabilities, which might require constant psychosocial and medical care, thus significantly reducing the quality of life. Some toxicants were identified as contributing factors in neurodevelopmental disorders (NDDs), such as autism spectrum disorder (ASD), intellectual disability (ID), attention deficit/hyperactivity disorder (ADHD), social communication disorders, as well as motor disorders (Hansen and Rogers, 2013). Currently approximately (approx.) 10–15% of all births are affected by one or a combination of some NDDs and this trend seems to be growing (Bloom et al., 2013). With the contribution of genetic predispositions estimated on maximum 40% of all cases, environmental exposures to toxins (alone or by interaction with genetic factors) are considered as a main cause of NDDs (Grandjean and Landrigan, 2014). Therefore, there is an urgent need to identify and further study these substances.

In developmental neurotoxicity research, the use of mammalian models is still a primary choice. However, the time and expense involved in analyzing a growing number of compounds have recently led to the adaptation of an alternative *in vivo* and *in vitro* models, which provide high-throughput screens much faster and at lower cost (Peterson et al., 2008). This review discusses the benefits of using

* Corresponding author.

E-mail addresses: joanna.ruszkiewicz@einstein.yu.edu (J.A. Ruszkiewicz), mahfuzur.miah@phd.einstein.yu.edu (M.R. Miah), ayodele.akinyemi@einstein.yu.edu (A.J. Akinyemi), omamuyovwi.ijomone@einstein.yu.edu (O.M. Ijomone), michael.aschner@einstein.yu.edu (M. Aschner).

<https://doi.org/10.1016/j.taap.2018.03.016>

Received 17 October 2017; Received in revised form 28 February 2018; Accepted 12 March 2018

Available online 14 March 2018

0041-008X/ © 2018 Elsevier Inc. All rights reserved.

Caenorhabditis elegans (*C. elegans*) as an alternative and complementary model to mammalian systems and gives examples of the uses of *C. elegans* in neurodevelopmental toxicology research.

2. *C. elegans* in neuroscience

2.1. General remarks on *C. elegans* as model organism

The small non-parasitic nematode *C. elegans* is one of the best-established animal models that has contributed greatly to the understanding of many human diseases. It is a very attractive experimental model due to many advantages: small size – adults are approx. 1 mm in length, short life cycle (approx. 3 days at 20 °C), ability to self-fertilize and high reproductive rate (> 300 offspring per hermaphrodite), which makes its maintenance in the lab relatively easy and inexpensive. *C. elegans* lab assays are usually rapid, of low cost, and amenable to high-throughput analysis. Additionally, the worms' transparency and the ease of making reporter gene fusions (e.g. with green fluorescent protein, GFP) enable visualization of cell morphology and protein expression patterns *in vivo*. The worm is also a powerful genetic tool. Their genome is easy to manipulate - gene knockdowns (KD) can be generated with RNAi and knockouts (KO) can be easily generated by directed mutagenesis. Moreover, the self-fertilizing hermaphrodite allows for homozygosity of mutations and mating with males can produce strains with multiple mutations. The nematode was the first multicellular organism to have its genome fully sequenced (Consortium and T.C.e.S., 1998), revealing a high degree of evolutionary conservation with higher eukaryotes and mammals. Worms express homologues to approx. 80% of human genes, many of which share great sequence identity (Culetto and Sattelle, 2000; Kennedy, 2008). In addition, the basic biological functions and many biochemical pathways are highly conserved with higher organisms, such as apoptosis (Malin and Shaham, 2015), stress response (Rodriguez et al., 2013) and various cell signaling pathways (Blaxter, 2011; Lapierre and Hansen, 2012; Sato et al., 2014), allowing for findings from *C. elegans* to be extrapolated and further confirmed in vertebrate systems.

C. elegans, however, poses several limitations as a model organism, which should not be omitted, when discussing its utility. This simple organism lacks many specific organs, such as kidneys, liver, lung, skin and a circulatory system. The evolutionary distance from humans might be a limitation, especially when studying complex conditions. Self-fertilization and the scarcity of males in wild type (WT) population make it difficult to study sex-specific differences, an approach currently promoted in neuroscience, including developmental neurotoxicology, and in other fields (Brooks and Clayton, 2017).

Nevertheless, the numerous advantages highly outrank these limitations, and the worms have been successfully utilized as model organism in many studies.

2.2. The nervous system of *C. elegans*

One of the biggest advantages of the model is its nervous system, which is well characterized and considered as structurally and functionally similar to mammals. The nematode nervous system contains 302 neurons in adult hermaphrodites (383 in males), divided into 118 morphologically distinct classes and 56 glia cells, which together form over 7600 synapses (White et al., 1986). *C. elegans* is the only organism whose neuronal wiring diagram has been determined – all neurons and the major interactions between them are well characterized and mapped. The biochemistry of the *C. elegans* nervous system is also highly conserved with mammals – as the worms express similar ion channels, receptors, vesicular transporters, and other synaptic components (Bargmann, 1998). Furthermore, formation, trafficking and release of synaptic vesicles are analogous to mammals (Barclay et al., 2012). The nematode uses classical neurotransmitters such as glutamate (Glu), γ -aminobutyric acid (GABA), dopamine (DA), serotonin (5-

hydroxytryptamine, 5-HT) and acetylcholine (ACh) (Brownlee and Fairweather, 1999). However, they seem to lack epinephrine, nor-epinephrine and histamine signaling, and some other significant differences have been described in sodium-dependent channels (Bargmann, 1998; Goodman et al., 1998). Apart from the cellular and molecular similarity to higher organism, *C. elegans* exhibit variety of behaviors, including social interaction, which might contribute to the understanding of the nervous system function. Because of its simplicity, investigating worms' behavior may be free from confounding factors, present in higher animals (Williams and Dusenbery, 1990).

2.3. *C. elegans* in neurotoxicology

Structural and functional similarities to the human nervous system, together with other advantages, strengthen the worms' choice as a model in neuroscience. Worms' transparency allows the visualization of neurons and synapses *in vivo* by fluorescent markers (Chalfie et al., 1994), and neuronal excitability can be followed by live calcium imaging (Nguyen et al., 2016), which allows inference on neuronal activity with behavioral phenotypes. Moreover, its ease of genetic manipulations allows the identification of genes significant for neuronal formation, migration, activity, or other functions. Since Sydney Brenner's Nobel Prize-winning studies, using *C. elegans* to investigate development and function of nervous system (Brenner, 1973), the nematode contributed greatly to understanding brain physiology. For instance, worms played a great role in determining the role of synaptic proteins such as neuroligin and neurexin in ASD (Schmeisser and Parker, 2018) and used in studies of neurodegenerative diseases, like Alzheimer's disease (AD) and Parkinson's disease (PD), shedding light on their causes and progression (Calahorra and Ruiz-Rubio, 2011).

The field of neurotoxicology has also taken advantage of the model. The use of *C. elegans* in toxicological studies has recently escalated, mostly due to the ease of high-throughput chemical-genetic screens, and the ability to analyze the effect of toxic mixtures. Chronic and delayed effects of environmental toxicants are often difficult to monitor in mammalian models due to their long life cycles, therefore the short lifespan of *C. elegans* affords a convenient alternative in assessing this impact (Tejeda-Benitez and Olivero-Verbel, 2016; Honnen, 2017; Hunt, 2017). To date, almost every type of known toxicant has been tested in this animal model (Tejeda-Benitez and Olivero-Verbel, 2016; Hunt, 2017). Many toxicants, including metals and pesticides, have been examined with respect to their effects on the *C. elegans* nervous system and very frequently results from these studies have closely emulated those observed in mammalian systems.

This review focuses specifically on neurodevelopment and aims to highlight the contribution and utility of *C. elegans* to the field of neurodevelopmental toxicology. This review gives examples of the uses of *C. elegans* in neurodevelopmental toxicology research, summarized in Table 1, and discusses relevance to mammalian studies.

3. Neurodevelopmental toxicity studies in *C. elegans*

3.1. Manganese

Manganese (Mn) is one of the most plentiful naturally occurring trace elements. In organisms it exists in all tissues, where at physiological levels it regulates various cellular processes including metabolism of proteins, lipids and carbohydrates, immune system, growth and development. However, at high concentrations, Mn can be toxic, particularly to the brain (Sidoryk-Wegrzynowicz and Aschner, 2013; Chen et al., 2015a; Peres et al., 2016). Excessive exposure to Mn may lead to its deposition and accumulation within the brain as it readily crosses the blood brain barrier (BBB). Particularly, dopaminergic (DAergic) neurons of the nigrostriatal region are compromised leading to a parkinsonian syndrome referred to manganism. The syndrome presents clinical indicators that are similar to PD (Chen et al., 2015a). Mn

Table 1
C.elegans as a model organism in neurodevelopmental toxicity studies.

Compound	Experimental design	Effect	Similar effects in mammals	Reference	
Manganese (Mn)	L1 0.5 h 50 mM in dH ₂ O	Increased DAergic degeneration, partially dependent on expression of SMF-1.	Yes	Settivari et al. (2009)	
	L1 0.5 h 0.001–100 mM in 85 mM NaCl	Increased DAergic, but not cholinergic, GABAergic degeneration; effect dependent on DAT-1, DOPs and SMF-1 and inhibited by SKN-1 overexpression.	Yes	Benedetto et al. (2010)	
	L1 0.5 h 50 mM in dH ₂ O	Increased DAergic degeneration; dependent on GST-1 and SKN-1, but not DAT-1 expression; the effect inhibited by KD of <i>jnk-1</i> , <i>ced-3</i> and <i>csp-1</i> .	Yes	Settivari et al. (2013)	
	L1 1 h 10, 25 mM in 85 mM NaCl	Increased DAergic degeneration and reduced basal slowing response. SLC30A10 overexpression attenuated DAergic injury, while KO worsened the effects of Mn.	Yes	Leyva-Illades et al. (2014)	
	L1 0.5 h 50 mM in 85 mM NaCl	Increased dauer movement due to <i>djr-1.2</i> -dependent disruption of DAergic signaling.	ND	Chen et al. (2015c)	
	L1 1 h 5, 10 mM in 85 mM NaCl	Increased DAergic degeneration and reduced basal slowing response; mutation in <i>trt-1</i> resisted compromised DAergic systems.	Yes	Ijomone et al. (2016)	
	L1 0.5 h	Increased DAergic degeneration; effect aggravated by loss of <i>hsp-1</i> .	Yes	Avila et al. (2016)	
	Mercury (Hg)	L1, L4 0.5, 6, 15 h 0.1–10 mM MeHgCl in M9	Unchanged DA and GABA neurons morphology.	No	Helmcke et al. (2009)
		L1 0.5 h 20 μM MeHgCl in M9	Degeneration of DAergic neurons later in life (96 h), increased ROS levels, upregulation of SKN-1.	Yes	Martinez-Finley et al. (2013a)
		L1 0.5 h 10, 20 μM MeHgCl in M9	Decreased DA-mediated locomotor activity.	Yes	Martinez-Finley et al. (2013b)
		L4 48, 72, 96 h 0.5–125 μM MeHgCl in NGM plates	Degeneration of DA neurons, increased ROS levels and GST mRNA, developmental defects, protective role of SKN-1.	Yes	Vanduyn et al. (2010)
		L1, L4 5, 48 h 0.3–200 μM HgCl ₂ 0.03–50 μM MeHgCl in K medium	MeHgCl-induced neurotoxic effects at concentrations lower than for HgCl ₂ .	Yes	McElwee and Freedman (2011)
		L2, L4 12 h 50, 100, 200 μM MeHgCl in K medium	Neurobehavioral defects partially prevented by preconditioning with mild heat shock.	ND	Ye et al. (2010)
L4 6 h 2.5–50 μM HgCl ₂		Decreased associative learning behavior, thermotaxis and locomotion behavior.	Yes	Zhang et al. (2010)	
L1–L4 6 h 2.5–50 μM HgCl ₂ in K medium		Degeneration of GABAergic neurons.	Yes	Xing et al. (2009)	
Arsenic (As)		L4 24 h 100 μM As (III) in K-medium	Deficits in AFD sensory neurons, decrease in isothermal tracking behavior, altered locomotor behaviors.	Yes	Yu and Liao (2014)
		Egg to adult chronic exposure 1 mM As (III) in NGM plates	Increased sensitivity in paralysis assays suggesting alterations in cholinergic system; the role of <i>abts-1</i> in this process.	Yes	Liao et al. (2010)
Lead (Pb)	L1 worms pre-treated with 0.01 μM Se(IV) 40 h	Decreased locomotor behaviors: body bends, head thrashing, and reversal frequency; deficits in AFD sensory neurons; effects mitigated by Se pre-treatment.	Yes	Li et al. (2013)	
	L4 24 h 100 μM Pb(NO ₃) ₂ in K-medium	Decreased locomotor behaviors: body bends and head thrashes.	Yes	Sun et al. (2016)	
	L1–young adult 2.5 days 1.45 mg/l Pb in K-medium	Impaired association learning behavior.	Yes	Zhang et al. (2010)	
	L4 6 h 2.5, 10 μM Pb(NO ₃) ₂ in K-medium	Dose-dependent increase in dorsal and ventral cord gaps and increased loss of neuronal cell bodies in L1–L3 worms; deficits in cholinergic transmission in L1–L4 worms.	Yes	Xing et al. (2009)	
	L1–L4 6 h 2.5, 50, 100 μM Pb(NO ₃) ₂ in K-medium				
	Sodium fluoride (NaF)	L4 24 h 0.038, 0.38, 3.8 mM in M9	Defects in locomotion behavior, increased ROS levels and upregulation of oxidative stress-related genes.	Yes	Li et al. (2012)

(continued on next page)

Table 1 (continued)

Compound	Experimental design	Effect	Similar effects in mammals	Reference
Dichlorvos	L4 2, 8, 26 h 6, 15 µM in M9	Altered gene expression of several neuronal growth/repair-related genes.	Yes	Lewis et al. (2013)
Monocrotophos	L4 24 h 0.5, 0.75, 1.5 mM in K medium	Decreased AChE, enhancing effects for a high glucose diet.	Yes	Salim and Rajini (2014)
TouchDown/mancozed	L2 0.5 h 0.22–12.5% in M9	Neurodegeneration of DA and GABA systems.	ND	Negga et al. (2011), Negga et al., (2012)
EPTC/molinate/ MeDETC	L1 1 h 0.2, 0.4, 0.1 mM in M9	Changed DAergic cell morphology, decreased DA content and impaired DA-related behavior.	ND	Caito et al. (2013a)

Abbreviations: *abts-1*: bicarbonate transporter; ACh: acetylcholine; AChE: acetylcholinesterase; *ced-3*: cell death protein 3 subunit 2; *csp-1*: caspase A subunit p16; DA: dopamine; DAT-1: sodium-dependent dopamine transporter; dH₂O: distilled water, *djr-1.2*: glutathione-independent glyoxalase; DMT-1 divalent metal transporter 1; DOPs: dopamine receptors; EPTC: S-ethyl N,N-dipropylthiocarbamate; ERM: medaka embryo-rearing medium; GABA: γ-aminobutyric acid; GST: glutathione transferase; Hg: mercury; *hsp-1*: heat shock protein A; *jnk-1*: stress-activated protein kinase; M9: M9 minimal medium; MeDETC: S-methyl-N,N-diethylthiocarbamate; MeHgCl: methylmercury chloride; Mn: manganese; mRNA: messenger ribonucleic acid; ND: no data; NGM: nematode growth medium; NPs: nanoparticles; ROS: reactive oxygen species; Se: selenium; SKN-1: homologue of nuclear factor (erythroid-derived 2)-like 2; SLC30A10: solute carrier family 30 member 10; SMF-1: NRAMP-like transporter; *trt-1*: telomerase reverse transcriptase.

exposure may also present a risk factor for the development of PD (Guilarte, 2013).

Several studies have reported toxic effects of Mn exposure on children in relation to neurodevelopment (Coetzee et al., 2016; Bjorklund et al., 2017). For example, a study of 7–9-year-old children residing near a ferromanganese industry showed that both high and low concentrations of Mn in blood and hair are associated with lower intelligence quotient (IQ) scores (Haynes et al., 2015). Increased prenatal and early childhood exposure to Mn in drinking water has been shown to elevate the risk of neurobehavioral problems of children at 10 years of age (Rahman et al., 2017). Animal studies have also demonstrated developmental neurotoxicity associated with Mn exposure (Kikuchi et al., 2015; Peres et al., 2015), however the molecular mechanisms by which Mn imparts neurodevelopment remains unresolved, and in these regards, the use of *C. elegans* in Mn neurotoxicity is applied (Chen et al., 2015b).

Several studies from our lab and others have shown that *C. elegans* treatment with Mn at early life (Larva-1 stage, L1) induces DAergic neurodegeneration in L1, L4 and young adults (Settivari et al., 2009; Benedetto et al., 2010; Settivari et al., 2013; Leyva-Illades et al., 2014; Avila et al., 2016; Ijomone et al., 2016). Additionally, treatment of L1 worms with Mn altered neurobehavioral parameters at a later stage. Mn treatment at L1 increased dauer movement of WT worms at dauer stage (Chen et al., 2015c) and reduced basal slowing response at L4 and young adult stage (Leyva-Illades et al., 2014; Ijomone et al., 2016). Increased dauer movement and reduced basal slowing response are indicative of compromised DAergic neurotransmitter systems. Benedetto et al. (2010) suggested that Mn developmental neurotoxicity is specific to DAergic neurons in *C. elegans*. The authors did not observe any degeneration in cholinergic, GABAergic and chemosensory neurons, at doses that induced DAergic degeneration (Benedetto et al., 2010). However, it is possible that DAergic neurons are most sensitive to Mn developmental neurotoxicity, and higher exposure concentrations and duration may produce analogous degenerating effects in other neurotransmitter systems in *C. elegans*.

Similarly to higher organisms, in worms Mn has been reported to trigger reactive oxygen species (ROS) generation and induced oxidative stress (Settivari et al., 2009; Benedetto et al., 2010; Settivari et al., 2013; Avila et al., 2016), as well as alter activities of apoptotic proteins (Settivari et al., 2013) and PD-associated genes (Chen et al., 2015b). Expression of GST-1 (glutathione transferase 1) and SKN-1, a homologue of human nuclear factor (erythroid-derived 2)-like 2, NRF-2 – a transcription factor regulating antioxidant response involved in antioxidant and detoxification enzymes regulation, has been shown to

inhibit Mn-induced developmental neurotoxicity in *C. elegans* (Settivari et al., 2009; Benedetto et al., 2010; Settivari et al., 2013).

Involvement of apoptotic proteins has also been implicated in Mn-induced developmental neurotoxicity. Genetic KD mutations of *jnk-1*, *ced-3*, and *csp-1*, all inhibited DAergic neurodegeneration in young adult following exposure of L1 worms to Mn. JNKs belong to a class of MAPKs that are triggered by internal or external stressors, and are involved in proapoptotic signaling. CED-3 is homologous to vertebrate proapoptotic caspase-3, while CSP-1 is among three caspase-like proteins (the other two being CSP-2, CSP-3) that show some homology to vertebrate caspase-1-3, and may act as an initiator of apoptosis by upregulating CED-3 (Settivari et al., 2013).

Several transporters have been implicated in Mn-induced developmental neurotoxicity in mammals (Erikson et al., 2005; Au et al., 2008; Tuschl et al., 2012) and studies in worms have helped to understand their role. DAergic degeneration following Mn exposure in early stage of *C. elegans* development has been shown to be dependent on dopamine transporter DAT-1, as deletion mutations of *dat-1* protect against DAergic degeneration following Mn treatment. Nevertheless, *dat-1* mutants as well as mutants with KO of all DA receptors (*dop-1*, *dop-2*, *dop-3*) show overall hypersensitivity to Mn toxicity (decreased survival rate compared to WT controls). In addition, *C. elegans* with tyrosine hydroxylase (*cat-2*) and vesicular monoamine transporter (*cat-1*) loss of function mutations exhibit hyper-resistant to Mn treatment. DAT-1 and DOPs are associated with removal of DA from synapses, while internal DA generation and packaging for release at synapses depends on CAT-2 and CAT-1 respectively. Hence these results suggest extracellular, but not intracellular, DA exacerbates Mn neurotoxicity (Benedetto et al., 2010; Chen et al., 2015b). Contrastingly, Settivari et al. (2013) showed that Mn-induced DAergic degeneration is not dependent on DAT-1. These authors suggest that differences in the genetic background of the *dat-1* KO strain and WT strain used in the previous study (Benedetto et al., 2010) may have been the likely reason for this contrasting result. Developmental DAergic degeneration has been shown to be partially dependent on divalent metal transporter DMT-1 homologues SMF-1 in DAergic neurons, because *smf-1* deletion mutations partially protected against DAergic degeneration in worms (Benedetto et al., 2010; Settivari et al., 2013). Additionally, SLC30A10, an Mn efflux transporter has been shown to be involved in developmental neurotoxicity on DAergic system. Overexpression of SLC30A10 in worms reduced cellular Mn levels, thereby attenuating against Mn toxicity. Moreover, it significantly attenuated Mn-induced DAergic degeneration in L1 worms, while L89P mutation of SLC30A10 significantly worsened DAergic degeneration upon Mn exposure (Leyva-Illades et al., 2014).

Recent studies identified protein targets that may be involved in developmental neurotoxicity following Mn exposure. They include lipocalin-related protein LPR-5 (Rudgalvyte et al., 2016), telomerase reverse transcriptase TRT-1 (Ijomone et al., 2016), and heat shock proteins HSP-70, HSP-3 and CHN-1 (Avila et al., 2016). *lpr-5* KD resulted in increased animal death in both controls and Mn exposed L1 worms. Although the function of LPR-5 is not clear, it is suggested that it may be involved in Mn induced toxicity via endoplasmic reticulum-lipocalin-related pathways (Rudgalvyte et al., 2016). Mutation in TRT-1 (the catalytic subunit of telomerase) in *C. elegans* inhibited DAergic degeneration in L1 worms upon Mn exposure. However, it may be due to the ability of *C. elegans* to survive without functional telomerase (Ijomone et al., 2016). Loss of *hsp-70*, *hsp-3* and *chn-1* reduced survival rate as well as increased protein oxidation following Mn exposure to L1 worms. Particularly, loss of *hsp-70* worsened Mn-induced DAergic degeneration. This effect may involve the blocking of transcriptional up-regulation of *pink-1*, a gene linked to PD (Avila et al., 2016).

PD-associated genes, like *parkin*, *dj-1* and α -synuclein, have been implicated in Mn-induced DAergic degeneration (Chen et al., 2015a). Overexpression of α -synuclein in worms rescued DAergic degeneration in the background of *parkin* deletion mutation, but not *dj-1* mutation (Chakraborty et al., 2013). However, it should be noted that worms expressing human WT α -synuclein showed increased DAergic degeneration following Mn treatment reinforcing the exacerbating role exerted by α -synuclein in PD (Settivari et al., 2009). *djr-1.2* deletion increased dauer movement following Mn exposure, suggesting *djr-1.2* deletion disrupts DAergic signaling; this effect was exacerbated upon Mn exposure during *C. elegans* development (Chen et al., 2015c).

In summary, *C. elegans*' unique properties contributed greatly to better understanding of mechanism of Mn-induced developmental neurotoxicity. Worms' transparency and ability to visualize neurons allowed to identify DAergic system as a key target for neurodegeneration upon early life Mn exposure (Settivari et al., 2009; Benedetto et al., 2010; Settivari et al., 2013; Leyva-Illades et al., 2014; Avila et al., 2016; Ijomone et al., 2016), which was further confirmed by numerous behavioral tests (Chen et al., 2015c; Leyva-Illades et al., 2014; Ijomone et al., 2016). Moreover, ease of genetic manipulation allowed deeper insight into the mechanism, revealing the important role of extracellular DA and DA transporters in DAergic neurodegeneration (Benedetto et al., 2010; Chen et al., 2015b). Furthermore, genetic modifications allowed to reveal the significance of several antioxidant and detoxification genes (Settivari et al., 2009; Benedetto et al., 2010; Settivari et al., 2013; Avila et al., 2016) and heat-shock proteins (Avila et al., 2016) in Mn resistance, as well as the role of apoptotic proteins (Settivari et al., 2013), metal transporters (Benedetto et al., 2010; Settivari et al., 2013), or PD-associated genes (Chakraborty et al., 2013; Chen et al., 2015c) in the mechanism of Mn-induced neurotoxicity and neurodegeneration during early development.

3.2. Mercury

Mercury (Hg), especially its organic form – methylmercury (MeHg) is one of the most poisonous neurotoxicants found in the environment. Humans are exposed to MeHg mainly through consumption of contaminated fish and plants, but also through occupational exposure and anthropogenic mercury emissions (Clarkson and Magos, 2006; Rice et al., 2014). Acute exposure to MeHg has historically been seen by incidents in Minamata Bay (Japan) and in Barsa (Iraq). Even though over the past few years MeHg exposure has been substantially curbed, one is still affected because it is so persistent in the environment, posing a risk related to low-levels chronic exposures (Grandjean et al., 2010).

MeHg crosses the blood-brain and placental barriers, causing a serious threat to developing brains in fetuses and infants. The neurodevelopmental toxicity of MeHg is well established -numerous studies addressed MeHg developmental neurotoxicity in animals and humans, and linked it to ROS accumulation, oxidative stress, disruption in

neurotransmitters or metallothioneins homeostasis (Sanfeliu et al., 2003; Bose-O'Reilly et al., 2010; Llop et al., 2013; Antonelli et al., 2017). However, the mechanism of toxicity is still not fully understood (Llop et al., 2015), and a lack of efficient treatment stimulate further research, in which *C. elegans* has already played a significant role.

To date, several initial studies aimed to characterize the effect of Hg and MeHg on the nematode in general, as well as on its developing nervous system, revealing numerous similarities with mammalian models. MeHg has been shown to be transported in the worm by a conserved mechanism, dependent on amino acid transporters LAT1 (Caio et al., 2013b). MeHg was more toxic to L1 and L4 stage *C. elegans* than HgCl₂, when assessing feeding, movement, and reproduction, all of which require neuromuscular activity (McElwee and Freedman, 2011), and these results are consistent with studies in other species. When L4 larval were exposed to a low (2.5 μ M) HgCl₂ levels for 6 h, it caused significant decrease of associative learning behavior, whereas 50 μ M caused significant reduction of body bend frequency and thermotaxis to cultivation temperature (Zhang et al., 2010). One of the first studies showed that L1 worms are significantly more sensitive than L4 to acute MeHgCl treatment, likely due to undeveloped detoxification system (Helmcke et al., 2009). Moreover, MeHg treatment inhibited worms' development and pharyngeal pumping, however DAergic and GABAergic neurons morphology was not impaired in animals that survived MeHgCl exposure (up to 72 h post-treatment), suggesting that *C. elegans* neurons may be less sensitive than mammalian, even though Hg levels found in *C. elegans* were similar to those causing deleterious alterations in mammalian systems (Helmcke et al., 2009). Degeneration of GABAergic neurons has been observed after treatment with HgCl₂ (up to 100 μ M, 6 h); of importance, younger larvae (L1–L3) showed greater sensitivity to neurotoxicity (assessed by changes in neuronal survival and synaptic function) than L4 larvae and young adult nematodes (Xing et al., 2009). Martinez-Finley et al. has reported a loss of DAergic neurons later in life (96 h) following early-life (L1) MeHg exposure (0.5 h, 20 μ M). Moreover, loss-of-function of DAergic neurons (basal slowing response) was observed later in life (72 h) (Martinez-Finley et al., 2013a). The role of SKN-1 was investigated in this context, showing that DA levels were decreased in *skn-1* KO compared to WT worms, following early life exposure to MeHg. ROS levels were elevated in *skn-1* KO and in WT worms following MeHg exposure. SKN-1 expression in the gut has been seen 24 h following exposure to 20 μ M MeHg (Martinez-Finley et al., 2013a). The upregulation of NRF-2 and genes under its control has been shown in many mammalian models upon exposure to MeHg (Ni et al., 2010; Kumagai et al., 2013; Culbreth et al., 2017) and other developmental neurotoxicants (Lau et al., 2013). A previous study has also shown that reduction in SKN-1 gene expression increases L4 worms vulnerability to MeHg, affects expression of GST, and prevents DAergic degeneration.

The role of mammalian parkin/PARK2 protein was also studied in MeHg neurotoxicity using *C. elegans* model. The *pdr-1* KO exacerbated MeHg toxicity and damage to the DAergic system, however a change in DA-mediated locomotor activity (basal slowing response) was observed in WT worms, but not in *pdr-1* KO (Martinez-Finley et al., 2013b). The role of metallothioneins in adaptation response to Hg has been also investigated in *C. elegans*. Pre-treatment with mild heat-shock (1 h at 36 °C) at L2 larva stage prevented neurobehavioral defects and the activation of stress response in nematodes exposed (L4, 12 h) to mercury at lower concentrations (50 and 100 μ M), but not at higher levels (200 μ M) (Ye et al., 2010).

To date, numerous neurobehavioral studies have confirmed MeHg-induced neurodevelopmental toxicity in young *C. elegans* (Helmcke et al., 2009; Martinez-Finley et al., 2013a; McElwee and Freedman, 2011; Zhang et al., 2010), while neuronal visualization revealed DAergic and GABAergic neurodegeneration later in life (Helmcke et al., 2009; Martinez-Finley et al., 2013a; Xing et al., 2009). Furthermore, genetic engineering allowed better understanding of the protective role of *skn-1* (Martinez-Finley et al., 2013a) and metallothioneins (Ye et al.,

2010) in MeHg resistance, as well as the contribution of *pdr-1* (Martinez-Finley et al., 2013b) in the mechanisms of MeHg-induced developmental neurotoxicity.

3.3. Arsenic

Arsenic (As) and arsenic compounds are known carcinogens that are widely present in the environment and are particularly elevated in groundwater in a variety of nations around the world due to both natural and anthropogenic sources. The World Health Organization has set 10–50 ppb as provisional permissible concentration limits for As in groundwater, yet many areas around the world have populations that are exposed to ten to hundreds of times those limits (Shankar et al., 2014). General consequences of As exposure include, but are not limited to, arsenicosis, skin melanosis, cancers of various organs, and IQ declines (O'Bryant et al., 2011; Hunt et al., 2014; Tsuji et al., 2015). Though extensively studied in the context of cancers, much remains to be understood in terms of the long-term effects of early life arsenic exposure on neurodevelopment and how early life toxicant insults or low level chronic exposures may contribute to neurological diseases.

The role of *abts-1* (bicarbonate transporter) in As neurotoxicity was evaluated in *C. elegans*. The experimental model utilized WT and *abts-1* mutants grown on plates containing 1 mM As from egg stage to adult (approx. 2.5 days). The *abts-1* expression was increased in neurons of transgenic worms expressing *Pabts-1::GFP* growing on As-containing plates. As-treated worms showed increased sensitivity in a paralysis assay performed with exposure to levamisole and aldicarb - the finding alludes to the involvement of *abts-1* and arsenic toxicity in alterations of the cholinergic system (Liao et al., 2010).

When L4 worms were treated with As (10–200 μ M) for 24 h, it resulted in robust changes in a variety of locomotor behaviors. Worms tagged with GFP in AFD thermosensory neurons showed decreases in cell body size and fluorescence intensity when comparing 100 μ M As treated worms to control (Yu and Liao, 2014). The AFD neurons are two ciliated neurons responsible for a variety of functions related to thermotactic behaviors and interacts with the AIY interneuron (Clark et al., 2006). These visible changes in the neurons also reflected functional change as As-treated worms exhibited a significant decrease in isothermal tracking (IT) behavior. Though the paper focuses on neurotoxic effects in AFD neurons, the locomotor behavior deficits suggest impairments of other neurons. Changes in the body bends and head thrashing allude to possible direct changes to motor neurons; loss of the AIZ interneuron has been found to lead to hyporeversal effects, similar to what was observed in worms treated with As, suggesting another neuron might be affected (Tsalik and Hobert, 2003). Reactive oxygen species (ROS) levels were also increased in L4 worms treated with As, and the As-induced toxicity effects were ameliorated in a variety of neurobehavioral assays when worms were pretreated with curcumin, a natural antioxidant, further highlighting that the mechanism of neurotoxicity involves oxidative processes (Yu and Liao, 2014).

Other literature utilizing the *C. elegans* model for As research has built on the observation of increased ROS levels post-As treatment at various life stages of the worm and connected it to acceleration of aging and mitochondrial dysfunction (Luz et al., 2016; Yu et al., 2016; Luz et al., 2017). These findings were whole body effects and were not evaluated specifically in neurons, and did not take advantage of potentially exposing worms at earlier life stages in development that could have given further insight into the effects of chronic arsenic exposure on neurodevelopment. Sahu et al. (2013) treated early development L1 stage worms with As, and although they did not look directly at neuronal changes, they did examine changes in gene expression - of particular interest were changes in ferroportin (responsible for iron export out of the cell) and zinc ion binding gene classes after As treatment (Sahu et al., 2013). Changes in iron and zinc homeostasis have been linked to a variety of neurological diseases including, but not limited to, AD, PD, ADHD, amyotrophic lateral sclerosis (ALS), and depression.

Thus, results from this paper suggest that early-life As exposure may contribute to neurological disease development by affecting the homeostasis of other biologically pertinent metals in the body (Hare et al., 2013; Szewczyk, 2013).

To date, early life exposure to As in *C. elegans* has been linked to increased ROS production and neurodegeneration, especially of AFD neurons, followed by neurobehavioral changes, like impaired thermosensory function (Yu and Liao, 2014). Moreover, the involvement of *abts-1* in As developmental neurotoxicity has been demonstrated (Liao et al., 2010). These studies employed a key advantages of worm model - ease of neuronal visualization, conduction of behavioral assays and genetic manipulations.

3.4. Lead

Lead (Pb) is a naturally occurring heavy metal in the Earth's crust and a non-essential metal in human biology. Human exposure to Pb is mostly due to its industrial use in a variety of alloys and compounds. Lead and Pb-alloys can be found in radiation shielding, pipes, weights, ammunition and batteries, moreover, Pb compounds can be found in dyes, paints, and gasoline, though the amounts have been greatly reduced or eliminated in recent years with increased awareness of the detrimental effects of Pb poisoning and exposure. These effects include kidney function impairment, thyroid function impairment, cardiovascular disease, decreased reproductive health, encephalopathy, neurodevelopmental defects in children, preterm birth, and more. Delving into neurodevelopmental effects of Pb exposure, the literature notes changes and impairment in IQ, memory, attention, language comprehension, processing speed, motor function and affect (Abadin et al., 2007; Mason et al., 2014). Molecular mechanism of Pb neurotoxicity is complex and includes oxidative stress, membrane disruptions, deregulation of cell signaling and impaired neurotransmission (Sanders et al., 2009).

The *C. elegans* model has been extensively used to examine Pb toxicity in general, but has not been specifically taken advantage of in the field of neurotoxicity and neurodevelopment, as noted by the scarcity of literature in this subfield. In one study, worms were pretreated with selenium (0.01 μ M Se(IV)) for 40 h starting from L1, and L4 worms were then exposed to 100 μ M Pb(NO₃)₂ for 24 h. The study showed decreased number of body bends, head thrashing, and reversal frequency after Pb exposure, which might be mitigated by pretreatment with Se (Li et al., 2013). Changes in locomotor assays could be attributed to changes in AFD neuron function, as stated before. Indeed, morphological changes in AFD neurons were observed after Pb-treatment, namely decreased cell body size. Moreover, a significant reduction in mRNA levels of *tax-2*, *tax-4*, and *ceh-14* were also observed after Pb exposure; these genes are required for proper AFD differentiation and function (Li et al., 2013). Decreased locomotor activity (body bends and head thrashes) was also shown when L1 worms were chronically treated with 1.45 mg/l Pb until young adult life stage (Sun et al., 2016).

When L4 worms were treated with 2.5 or 10 μ M Pb(NO₃)₂ for 6 h, they showed significant decrease in association learning behavior, as compared to control in a thermotaxis assay paradigm (Zhang et al., 2010). In another study, the same group utilized L1–L4 worms expressing GFP in GABAergic neurons and treated them with Pb(NO₃)₂ solutions (2.5, 50, or 100 μ M) for 6 h. The study demonstrated that the Pb exposures led to significant dose-dependent increases in dorsal and ventral cord gaps and increases in loss of neuronal cell bodies for L1–L3 treated worms, reflecting axonal and cell body degeneration post-Pb treatment (Xing et al., 2009). Worms were also treated with aldicarb and levamisole with the same paradigm, to determine if Pb treatment impacted cholinergic neurons. Normally, worms are paralyzed in these assays and continued motility upon exposure to aldicarb or levamisole suggest pre- or post-synaptic changes in cholinergic neurons, respectively. Increased motility was observed for all three concentrations of Pb for L1–L3 worms, as well as for 50 and 100 μ M for L4 worms, and

100 μ M Pb treatment for young adult, suggesting deficits in cholinergic transmission. Throughout the paper, there is a consistent theme of increased maturity in life stage leading to greater resistance against the impacts of lead exposure (Xing et al., 2009). Again, major advantages of the model application in neuroscience – neuronal GFP labeling and relative simplicity of behavioral testing, have been employed to reveal the effects of Pb on *C. elegans* larvae, which were degeneration of AFD neurons (Li et al., 2013) and GABAergic neurons (Xing et al., 2009), as well as various neurobehavioral changes (Li et al., 2013; Sun et al., 2016; Xing et al., 2009; Zhang et al., 2010).

3.5. Fluoride

Fluoride (F) is an essential trace element ubiquitous in the environment (Whitford, 1983). Additionally, due to its positive effect on teeth and bones, one of fluoride compounds – sodium fluoride (NaF) – is supplemented worldwide in public drinking water. In 2012, 74.6% of people with access to public water systems, or nearly 211 million people in the United States, drank artificially fluoridated water (Chaudhary et al., 2010). Thus, human exposure to substantial amounts of this element occurs on regular basis. Although consumption of drinking water and usage of NaF-containing dental hygiene products is considered the major source of human intake, potential for bioaccumulation of fluoride ion in living organisms and environment results from additional sources of exposure, like consumption of seafood or plants grown in high-fluoride waters or soils (Ozsvath, 2009; Chaudhary et al., 2010). Despite its positive effects, some studies suggest that prolonged ingestion of fluoride produces deleterious effects, especially on the CNS, which poses serious global health concern (Gui et al., 2010). Sodium fluoride has been shown to induce neurological impairment particularly in the hippocampus and cerebral cortex in experimental animals (Bhatnagar et al., 2002; Sharma et al., 2014). Several epidemiological and experimental studies have highlighted closed relationship between fluoride ingestion and neurodevelopmental delays (Chioca et al., 2008; Choi et al., 2012), alike decreased IQ of children (Aravind et al., 2016). Previous studies have demonstrated that the toxic effect of fluoride exposure on the CNS could be attributed to their ability to cross the BBB and lead to oxidative stress, DNA damage and decrease in nicotinic acetylcholine receptors (Ranpariya et al., 2011; Sharma et al., 2014), however the mechanisms underlying its developmental neurotoxicity are not fully understood.

Recently, Li et al. reported that NaF exposure can induce multiple biological toxicities to *C. elegans* in a concentration-dependent manner at physiological, biochemical, and molecular levels. In this study *C. elegans* (L4 stage) were exposed to three concentrations of NaF (0.038, 0.38, 3.8 mM) for 24 h. Furthermore, NaF induced behavioral deficits – decreased head thrashes and body bend frequency, and increased ROS levels and expression of some oxidative stress-related genes: *hsp-16.1*, *sod-3*, *ctl-2*, *dhs-28*, *gst-1*, and *cep-1* (Li et al., 2012). These results are consistent with some observations in higher organisms (Dec et al., 2017).

3.6. Pesticides

Dichlorvos, an organophosphorus (OP) pesticide, is widely used globally due to its high potency and low price (Zhang et al., 2017). As an OP, it inhibits acetylcholinesterase (AChE) – the enzyme which removes acetylcholine (ACh) from the synapse – leading to an excess of ACh in the synapse and impaired nerve function (Kobayashi et al., 1986). Dichlorvos is classified by the World Health Organization (WHO) as highly hazardous and its detrimental effects on humans are likely related to continuous dietary uptake, including some nervous system diseases at relatively low exposure levels (Mackenzie Ross et al., 2010; Velmurugan et al., 2013; Du et al., 2014).

Lewis et al. studied the effects of developmental exposure to dichlorvos in the *C. elegans* model; following exposure of L4 worms to

dichlorvos, several neuronal growth/repair-related gene expressions were altered: *dlk-1* (MAPKKK) and *pmk-3* (p38 MAPK), genes related to axon regeneration, were upregulated; expression level of *unc-14*, *unc-129*, *eva-1*, *klc-2*, and *pak-1*, all related to axonal guidance, was also increased; finally, the components of the SMA TGF- β pathway – which is related to sensory neuronal development – were also upregulated. The suspected underlying mechanism for these findings is, according to the authors, mitochondrial dysfunction, which stems from the changes in metabolic processes found following dichlorvos exposure (Lewis et al., 2013). Similar observations were true for rodents (Masoud et al., 2009; Wani et al., 2011).

Monocrotophos (MCP), an OP insecticide, also works by inhibiting AChE; it is highly toxic, fast acting and non-organism specific, which is why its use is banned in most developed countries (Kavitha and Venkateswara Rao, 2007; Maniyar et al., 2011; Leelaja and Rajini, 2013). As a high glucose diet has already been associated with the exacerbation of OP toxicity (Olivier et al., 2001; Liu et al., 2007), Salim and Rajini (2014) used the *C. elegans* model to study the developmental neurotoxicity of MCP and the effects of a high-glucose diet on such toxicity. Following exposure of L4 worms to MCP, a dose-dependent decrease in AChE was reported for both control and high-glucose worm groups, with a lower mean activity in the high glucose group, suggesting an increased nervous-system vulnerability under the high glucose conditions.

Glyphosate, a glycine analogue, is the active ingredient in some of the most common herbicides globally; its popularity led to the development of genetically modified crops which are resistant to its effects (Woodburn, 2000; Gardner and Nelson, 2008; Dewar, 2009). Recent studies show that while glyphosate alone is relatively non-toxic, exposure to glyphosate-containing products is related to mitochondrial dysfunction (Bababunmi et al., 1979; Olorunsogo et al., 1979; Olorunsogo and Bababunmi, 1980; Peixoto, 2005). Mancozeb, a manganese (Mn) containing fungicide, acts by inhibiting enzymes and is massively used worldwide – approximately 3.6 million kilograms annually (University, 1987; Giannese and Marcelli, 2000; Gianessi and Reigner, 2006). Exposure to mancozeb-containing products has been associated with apoptosis, neurodegeneration and inhibition of mitochondrial respiration (Soleo et al., 1996; Calviello et al., 2006; Domico et al., 2006; Domico et al., 2007; Zhang et al., 2017). In order to study the potential developmental neurotoxicity of TouchDown (TD; a glyphosate-containing herbicide) and Mancozeb (MZ; a Mn-containing fungicide), the *C. elegans* model was used, specifically the NW1229 strain, which has all neurons tagged with GFP: L2 worms were exposed, acutely or chronically, to either product or both. Fluorescent microscopy analyses revealed dose-dependent neurodegeneration for all parameters, and to the highest extent following MZ exposure (Negga et al., 2011). In a follow-up study, using the same design, it was found that exposure to TD and MZ is associated with neurodegeneration of GABAergic and DAergic neurons, suggesting a more specific effect for these compounds in the nervous system (Negga et al., 2012). Further, impaired neuronal development and axon growth was observed in rat primary cultures of hippocampal pyramidal cells exposed to this herbicide (Coullery et al., 2016).

Thiocarbamate and dithiocarbamate pesticides are among the most commonly used pesticides in the US (Grube et al., 2011); their mechanism of action involves the metal-dependant and sulfhydryl enzyme systems in fungi, bacteria, plants, insects and mammals (Miller, 1982). A major concern regarding exposure to these compounds is their metabolites, which can lead to protein modification and enzyme inhibition associated with mammalian neurotoxicity (Savolainen and Hervonen, 1985; Pentylala and Chetty, 1993; Staub et al., 1995; Tonkin et al., 2004; Viquez et al., 2012). Caito et al. (2013a) studied the effects of S-ethyl N,N-dipropylthiocarbamate (EPTC), molinate, and S-methyl-N,N-diethylthiocarbamate (MeDETC) – two thiocarbamate pesticides and one reactive intermediate of their metabolism, respectively – on the development and function of the nervous system in *C. elegans*.

Following acute exposure during the L1 phase the following parameters were studied in the worms: neuronal morphology, neurotransmitter content and basal-slowness response behavior. Pesticide exposure was associated with a DAergic deficit, including loss of cell morphology, decreased DA content and impaired DA-related behavior. Similar findings were not found in other neurotransmitter systems (cholinergic, glutamatergic and GABAergic), suggesting a potential role these compounds may have in the development of PD.

4. Conclusions and perspectives

C. elegans is a model organism which allows to combine the study of neuronal development, connectivity, physiology and behavior, changed by environmental and genetic factors in a simplified, *in vivo* context. Numerous advantages favor its use in toxicology and make it a recent model of choice in studying neurodevelopmental disorders. From the reviewed data, it emerges that *C. elegans* can be a useful tool in evaluating the effects of toxins on developing nervous system, advancing our understanding of the molecular mechanisms underlying toxicity. With effects frequently analogous to those observed in mammals, there is a strong premise, that *C. elegans* findings can be extrapolated to higher organisms, including humans. Therefore, its experimental popularity will probably continue to grow and hopefully result in identification of new hazardous chemicals, as well as better insight into pathomechanisms of known neurotoxicants.

C. elegans has already been successfully implemented in general toxicity studies of other developmental neurotoxins, like ethanol (Davis et al., 2008) toluene (Davies et al., 2012), polychlorinated biphenyls (Menzel et al., 2007; Schafer et al., 2009), various flame retardants (Behl et al., 2016) and chlorpyrifos (Ruan et al., 2012), resulting in characteristic, pathological phenotypes; however, their neurodevelopmental effects in worms have yet to be addressed.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgment

This work has been supported by the National Institutes of Health to MA [grant numbers NIEHS R01ES07331, NIEHS R01ES10563 and NIEHS R01ES020852. OMI acknowledges the support of IBRO-ISN research fellowship.

References

Abadin, H., Ashizawa, A., Stevens, Y.W., Lladós, F., Diamond, G., Sage, G., Citra, M., Quinones, A., Bosch, S.J., Swarts, S.G., 2007. Agency for Toxic Substances and Disease Registry (ATSDR) Toxicological Profiles, Toxicological Profile for Lead. Agency for Toxic Substances and Disease Registry (US), Atlanta (GA).

Antonelli, M.C., Pallares, M.E., Ceccatelli, S., Spulber, S., 2017. Long-term consequences of prenatal stress and neurotoxicants exposure on neurodevelopment. *Prog. Neurobiol.* 155, 21–35.

Aravind, A., Dhanya, R.S., Narayan, A., Sam, G., Adarsh, V.J., Kiran, M., 2016. Effect of fluoridated water on intelligence in 10–12-year-old school children. *J. Int. Soc. Prev. Commun. Dent.* 6, S237–S242.

Au, C., Benedetto, A., Aschner, M., 2008. Manganese transport in eukaryotes: the role of DMT1. *Neurotoxicology* 29, 569–576.

Avila, D.S., Benedetto, A., Au, C., Bornhorst, J., Aschner, M., 2016. Involvement of heat shock proteins on Mn-induced toxicity in *Caenorhabditis elegans*. *BMC Pharmacol. Toxicol.* 17, 54.

Bababunmi, E.A., Olorunsogo, O.O., Bassir, O., 1979. The uncoupling effect of N-(phosphonomethyl)glycine on isolated rat liver mitochondria. *Biochem. Pharmacol.* 28, 925–927.

Barclay, J.W., Morgan, A., Burgoyne, R.D., 2012. Neurotransmitter release mechanisms studied in *Caenorhabditis elegans*. *Cell Calcium* 52, 289–295.

Bargmann, C.I., 1998. Neurobiology of the *Caenorhabditis elegans* genome. *Science (New York, N.Y.)* 282, 2028–2033.

Behl, M., Rice, J.R., Smith, M.V., Co, C.A., Bridge, M.F., Hsieh, J.H., Freedman, J.H., Boyd, W.A., 2016. Editor's highlight: comparative toxicity of organophosphate flame retardants and polybrominated diphenyl ethers to *Caenorhabditis elegans*. *Toxicol. Sci.* 154, 241–252.

Benedetto, A., Au, C., Avila, D.S., Milatovic, D., Aschner, M., 2010. Extracellular dopamine potentiates Mn-induced oxidative stress, lifespan reduction, and dopaminergic neurodegeneration in a BLI-3-dependent manner in *Caenorhabditis elegans*. *PLoS Genet.* 6.

Bhatnagar, M., Rao, P., Sushma, J., Bhatnagar, R., 2002. Neurotoxicity of fluoride: neurodegeneration in hippocampus of female mice. *Indian J. Exp. Biol.* 40, 546–554.

Bjorklund, G., Chartrand, M.S., Aaseth, J., 2017. Manganese exposure and neurotoxic effects in children. *Environ. Res.* 155, 380–384.

Blaxter, M., 2011. Nematodes: the worm and its relatives. *PLoS Biol.* 9, e1001050.

Bloom, B., Jones, L.I., Freeman, G., 2013. Summary health statistics for U.S. children: National Health Interview Survey, 2012. In: *Vital Health Stat. Ser. 10. Data National Health Surv.*, pp. 1–81.

Bose-O'Reilly, S., McCarty, K.M., Steckling, N., Lettmeier, B., 2010. Mercury exposure and children's health. *Curr. Probl. Pediatr. Adolesc. Health Care* 40, 186–215.

Brenner, S., 1973. The genetics of behaviour. *Br. Med. Bull.* 29, 269–271.

Brooks, C.E., Clayton, J.A., 2017. Sex/gender influences on the nervous system: basic steps toward clinical progress. *J. Neurosci. Res.* 95, 14–16.

Brownlee, D.J., Fairweather, I., 1999. Exploring the neurotransmitter labyrinth in nematodes. *Trends Neurosci.* 22, 16–24.

Caito, S.W., Valentine, W.M., Aschner, M., 2013a. Dopaminergic neurotoxicity of S-ethyl N,N-dipropylthiocarbamate (EPTC), molinate, and S-methyl-N,N-diethylthiocarbamate (MeDETC) in *Caenorhabditis elegans*. *J. Neurochem.* 127, 837–851.

Caito, S.W., Zhang, Y., Aschner, M., 2013b. Involvement of AAT transporters in methylmercury toxicity in *Caenorhabditis elegans*. *Biochem. Biophys. Res. Commun.* 435, 546–550.

Calahorra, F., Ruiz-Rubio, M., 2011. *Caenorhabditis elegans* as an experimental tool for the study of complex neurological diseases: Parkinson's disease, Alzheimer's disease and autism spectrum disorder. *Invertebr. Neurosci.* 11, 73–83.

Calviello, G., Piccioni, E., Boninsegna, A., Tedesco, B., Maggiano, N., Serini, S., Wolf, F.I., Palozza, P., 2006. DNA damage and apoptosis induction by the pesticide Mancozeb in rat cells: involvement of the oxidative mechanism. *Toxicol. Appl. Pharmacol.* 211, 87–96.

Chakraborty, S., Bornhorst, J., Nguyen, T.T., Aschner, M., 2013. Oxidative stress mechanisms underlying Parkinson's disease-associated neurodegeneration in *C. elegans*. *Int. J. Mol. Sci.* 14, 23103–23128.

Chalfie, M., Tu, Y., Euskirchen, G., Ward, W.W., Prasher, D.C., 1994. Green fluorescent protein as a marker for gene expression. *Science (New York, N.Y.)* 263, 802–805.

Chaudhary, V., Kumar, M., Sharma, M., Yadav, B.S., 2010. Fluoride, boron and nitrate toxicity in ground water of northwest Rajasthan, India. *Environ. Monit. Assess.* 161, 343–348.

Chen, P., Chakraborty, S., Mukhopadhyay, S., Lee, E., Paoliello, M.M., Bowman, A.B., Aschner, M., 2015a. Manganese homeostasis in the nervous system. *J. Neurochem.* 134, 601–610.

Chen, P., Chakraborty, S., Peres, T.V., Bowman, A.B., Aschner, M., 2015b. Manganese-induced neurotoxicity: from *C. elegans* to humans. *Toxicol. Res.* 4, 191–202.

Chen, P., DeWitt, M.R., Bornhorst, J., Soares, F.A., Mukhopadhyay, S., Bowman, A.B., Aschner, M., 2015c. Age- and manganese-dependent modulation of dopaminergic phenotypes in a *C. elegans* DJ-1 genetic model of Parkinson's disease. *Metallomics* 7, 289–298.

Chioca, L.R., Raupp, I.M., Da Cunha, C., Losso, E.M., Andreolini, R., 2008. Subchronic fluoride intake induces impairment in habituation and active avoidance tasks in rats. *Eur. J. Pharmacol.* 579, 196–201.

Choi, A.L., Sun, G., Zhang, Y., Grandjean, P., 2012. Developmental fluoride neurotoxicity: a systematic review and meta-analysis. *Environ. Health Perspect.* 120, 1362–1368.

Clark, D.A., Biron, D., Sengupta, P., Samuel, A.D., 2006. The AFD sensory neurons encode multiple functions underlying thermotactic behavior in *Caenorhabditis elegans*. *J. Neurosci.* 26, 7444–7451.

Clarkson, T.W., Magos, L., 2006. The toxicology of mercury and its chemical compounds. *Crit. Rev. Toxicol.* 36, 609–662.

Coetzee, D.J., McGovern, P.M., Rao, R., Harnack, L.J., Georgieff, M.K., Stepanov, I., 2016. Measuring the impact of manganese exposure on children's neurodevelopment: advances and research gaps in biomarker-based approaches. *Environ. Health* 15, 91.

Consortium, T.C.e.S., 1998. Genome sequence of the nematode. *C. elegans*: a platform for investigating biology. *Science* 282, 2012–2018.

Coullery, R.P., Ferrari, M.E., Rosso, S.B., 2016. Neuronal development and axon growth are altered by glyphosate through a WNT non-canonical signaling pathway. *Neurotoxicology* 52, 150–161.

Culbreth, M., Zhang, Z., Aschner, M., 2017. Methylmercury augments Nrf2 activity by downregulation of the Src family kinase Fyn. *Neurotoxicology* 62, 200–206.

Culetto, E., Sattelle, D.B., 2000. A role for *Caenorhabditis elegans* in understanding the function and interactions of human disease genes. *Hum. Mol. Genet.* 9, 869–877.

Davies, A.G., Friedberg, R.I., Gupta, H., Chan, C.L., Shelton, K.L., Bettinger, J.C., 2012. Different genes influence toluene- and ethanol-induced locomotor impairment in *C. elegans*. *Drug Alcohol Depend.* 122, 47–54.

Davis, J.R., Li, Y., Rankin, C.H., 2008. Effects of developmental exposure to ethanol on *Caenorhabditis elegans*. *Alcohol. Clin. Exp. Res.* 32, 853–867.

Dec, K., Lukomska, A., Maciejewska, D., Jakubczyk, K., Baranowska-Bosiacka, I., Chlubek, D., Wasik, A., Gutowska, I., 2017. The influence of fluoride on the disturbances of homeostasis in the central nervous system. *Biol. Trace Elem. Res.* 177, 224–234.

Dewar, A.M., 2009. Weed control in glyphosate-tolerant maize in Europe. *Pest Manag. Sci.* 65, 1047–1058.

Domico, L.M., Zeevalk, G.D., Bernard, L.P., Cooper, K.R., 2006. Acute neurotoxic effects of mancozeb and maneb in mesencephalic neuronal cultures are associated with mitochondrial dysfunction. *Neurotoxicology* 27, 816–825.

- Domico, L.M., Cooper, K.R., Bernard, L.P., Zeevalk, G.D., 2007. Reactive oxygen species generation by the ethylene-bis-dithiocarbamate (EBDC) fungicide mancozeb and its contribution to neuronal toxicity in mesencephalic cells. *Neurotoxicology* 28, 1079–1091.
- Du, L., Li, S., Qi, L., Hou, Y., Zeng, Y., Xu, W., Wang, H., Zhao, X., Sun, C., 2014. Metabonomic analysis of the joint toxic action of long-term low-level exposure to a mixture of four organophosphate pesticides in rat plasma. *Mol. BioSyst.* 10, 1153–1161.
- Erikson, K.M., John, C.E., Jones, S.R., Aschner, M., 2005. Manganese accumulation in striatum of mice exposed to toxic doses is dependent upon a functional dopamine transporter. *Environ. Toxicol. Pharmacol.* 20, 390–394.
- Gardner, J., Nelson, G.C., 2008. Herbicides, glyphosate resistance and acute mammalian toxicity: simulating an environmental effect of glyphosate-resistant weeds in the USA. *Pest Manag. Sci.* 64, 470–478.
- Gianessi, L., Reigner, N., 2006. Pesticide Use in U.S. Crop Production: 2002, With Comparison to 1992 & 1997: Fungicides & Herbicides Pesticide Use in US crop Production. CropLife Foundation, Washington, DC.
- Giannese, L., Marcelli, M., 2000. Pesticide use in US crop production: 1997. In: Agriculture. U.S.D.o..
- Goodman, M.B., Hall, D.H., Avery, L., Lockery, S.R., 1998. Active currents regulate sensitivity and dynamic range in *C. elegans* neurons. *Neuron* 20, 763–772.
- Grandjean, P., Landrigan, P.J., 2014. Neurobehavioural effects of developmental toxicity. *Lancet Neurol.* 13, 330–338.
- Grandjean, P., Satoh, H., Murata, K., Eto, K., 2010. Adverse effects of methylmercury: environmental health research implications. *Environ. Health Perspect.* 118, 1137–1145.
- Grube, A., Donaldson, D., Kiely, T., Wu, L., 2011. Pesticides Industry Sales and Usage 2006 and 2007 Market Estimates. U.S.E.P. Agency, Washington, D.C.
- Gui, C.Z., Ran, L.Y., Li, J.P., Guan, Z.Z., 2010. Changes of learning and memory ability and brain nicotinic receptors of rat offspring with coal burning fluorosis. *Neurotoxicol. Teratol.* 32, 536–541.
- Guillarte, T.R., 2013. Manganese neurotoxicity: new perspectives from behavioral, neuroimaging, and neuropathological studies in humans and non-human primates. *Front. Aging Neurosci.* 5, 23.
- Hansen, R.L., Rogers, S.J., 2013. Autism and Other Neurodevelopmental Disorders. American Psychiatric Association, Washington, DC.
- Hare, D., Ayton, S., Bush, A., Lei, P., 2013. A delicate balance: iron metabolism and diseases of the brain. *Front. Aging Neurosci.* 5, 34.
- Haynes, E.N., Sucharew, H., Kuhnell, P., Alden, J., Barnas, M., Wright, R.O., Parsons, P.J., Aldous, K.M., Praamsma, M.L., Beidler, C., Dietrich, K.N., 2015. Manganese exposure and neurocognitive outcomes in rural school-age children: the communities actively researching exposure study (Ohio, USA). *Environ. Health Perspect.* 123, 1066–1071.
- Helmcke, K.J., Syversen, T., Miller 3rd, D.M., Aschner, M., 2009. Characterization of the effects of methylmercury on *Caenorhabditis elegans*. *Toxicol. Appl. Pharmacol.* 240, 265–272.
- Honnan, S., 2017. *Caenorhabditis elegans* as a powerful alternative model organism to promote research in genetic toxicology and biomedicine. *Arch. Toxicol.* 91, 2029–2044.
- Hu, W.F., Chahrouh, M.H., Walsh, C.A., 2014. The diverse genetic landscape of neurodevelopmental disorders. *Annu. Rev. Genomics Hum. Genet.* 15, 195–213.
- Hunt, P.R., 2017. The *C. Elegans* model in toxicity testing. *J. Appl. Toxicol.* 37, 50–59.
- Hunt, K.M., Srivastava, R.K., Elms, C.A., Athar, M., 2014. The mechanistic basis of arsenicosis: pathogenesis of skin cancer. *Cancer Lett.* 354, 211–219.
- Ijomone, O.M., Miah, M.R., Peres, T.V., Nwoha, P.U., Aschner, M., 2016. Null allele mutants of trt-1, the catalytic subunit of telomerase in *Caenorhabditis elegans*, are less sensitive to Mn-induced toxicity and DAergic degeneration. *Neurotoxicology* 57, 54–60.
- Kavitha, P., Venkateswara Rao, J., 2007. Oxidative stress and locomotor behaviour response as biomarkers for assessing recovery status of mosquito fish, *Gambusia affinis* after lethal effect of an organophosphate pesticide, monocrotophos. *Pestic. Biochem. Physiol.* 87, 182–188.
- Kennedy, B.K., 2008. The genetics of ageing: insight from genome-wide approaches in invertebrate model organisms. *J. Intern. Med.* 263, 142–152.
- Kikuchi, Y., Abe, H., Tanaka, T., Kato, M., Wang, L., Ikarashi, Y., Yoshida, T., Shibutani, M., 2015. Relationship between brain accumulation of manganese and aberration of hippocampal adult neurogenesis after oral exposure to manganese chloride in mice. *Toxicology* 331, 24–34.
- Kobayashi, H., Yuyama, A., Chiba, K., 1986. Cholinergic system of brain tissue in rats poisoned with the organophosphate, 0,0-dimethyl 0-(2,2-dichlorovinyl) phosphate. *Toxicol. Appl. Pharmacol.* 82, 32–39.
- Kumagai, Y., Kanda, H., Shinkai, Y., Toyama, T., 2013. The role of the Keap1/Nrf2 pathway in the cellular response to methylmercury. *Oxidative Med. Cell. Longev.* 2013, 848279.
- Lapierre, L.R., Hansen, M., 2012. Lessons from *C. elegans*: signaling pathways for longevity. *Trends Endocrinol. Metab.* 23, 637–644.
- Lau, A., Whitman, S.A., Jaramillo, M.C., Zhang, D.D., 2013. Arsenic-mediated activation of the Nrf2-Keap1 antioxidant pathway. *J. Biochem. Mol. Toxicol.* 27, 99–105.
- Leelaja, B.C., Rajini, P.S., 2013. Biochemical and physiological responses in *Caenorhabditis elegans* exposed to sublethal concentrations of the organophosphorus insecticide, monocrotophos. *Ecotoxicol. Environ. Saf.* 94, 8–13.
- Lewis, J.A., Gehman, E.A., Baer, C.E., Jackson, D.A., 2013. Alterations in gene expression in *Caenorhabditis elegans* associated with organophosphate pesticide intoxication and recovery. *BMC Genomics* 14, 291.
- Leyva-Ilades, D., Chen, P., Zogzas, C.E., Hutchens, S., Mercado, J.M., Swaim, C.D., Morrisett, R.A., Bowman, A.B., Aschner, M., Mukhopadhyay, S., 2014. SLC30A10 is a cell surface-localized manganese efflux transporter, and parkinsonism-causing mutations block its intracellular trafficking and efflux activity. *J. Neurosci.* 34, 14079–14095.
- Li, Q., Zhang, S.H., Yu, Y.H., Wang, L.P., Guan, S.W., Li, P.F., 2012. Toxicity of sodium fluoride to *Caenorhabditis elegans*. *Biomed. Environ. Sci.* 25, 216–223.
- Li, W.H., Shi, Y.C., Tseng, I.L., Liao, V.H., 2013. Protective efficacy of selenite against lead-induced neurotoxicity in *Caenorhabditis elegans*. *PLoS One* 8, e62387.
- Liao, V.H., Liu, J.T., Li, W.H., Yu, C.W., Hsieh, Y.C., 2010. *Caenorhabditis elegans* bicarbonate transporter ABTS-1 is involved in arsenite toxicity and cholinergic signaling. *Chem. Res. Toxicol.* 23, 926–932.
- Liu, J., Gupta, R.C., Goad, J.T., Karanth, S., Pope, C., 2007. Modulation of parathion toxicity by glucose feeding: is nitric oxide involved? *Toxicol. Appl. Pharmacol.* 219, 106–113.
- Llop, S., Lopez-Espinosa, M.J., Rebagliato, M., Ballester, F., 2013. Gender differences in the neurotoxicity of metals in children. *Toxicology* 311, 3–12.
- Llop, S., Ballester, F., Broberg, K., 2015. Effect of gene-mercury interactions on mercury toxicokinetics and neurotoxicity. *Curr. Environ. Health Rep.* 2, 179–194.
- Luz, A.L., Godebo, T.R., Bhatt, D.P., Ilkayeva, O.R., Maurer, L.L., Hirschey, M.D., Meyer, J.N., 2016. From the cover: arsenite uncouples mitochondrial respiration and induces a Warburg-like effect in *Caenorhabditis elegans*. *Toxicol. Sci.* 152, 349–362.
- Luz, A.L., Godebo, T.R., Smith, L.L., Leuthner, T.C., Maurer, L.L., Meyer, J.N., 2017. Deficiencies in mitochondrial dynamics sensitize *Caenorhabditis elegans* to arsenite and other mitochondrial toxicants by reducing mitochondrial adaptability. *Toxicology* 387, 81–94.
- Mackenzie Ross, S.J., Brewin, C.R., Curran, H.V., Furlong, C.E., Abraham-Smith, K.M., Harrison, V., 2010. Neuropsychological and psychiatric functioning in sheep farmers exposed to low levels of organophosphate pesticides. *Neurotoxicol. Teratol.* 32, 452–459.
- Malin, J.Z., Shaham, S., 2015. Cell death in *C. Elegans* development. *Curr. Top. Dev. Biol.* 114, 1–42.
- Maniyar, R.A., Ahmed, R.N., David, M., 2011. Monocrotophos: toxicity evaluation and respiratory responses of *Cyprinus carpio* (Linnaeus). *Recent Res. Sci. Technol.* 3, 51–54.
- Martinez-Finley, E.J., Caito, S., Slaughter, J.C., Aschner, M., 2013a. The role of skn-1 in methylmercury-induced latent dopaminergic neurodegeneration. *Neurochem. Res.* 38, 2650–2660.
- Martinez-Finley, E.J., Chakraborty, S., Slaughter, J.C., Aschner, M., 2013b. Early-life exposure to methylmercury in wildtype and pdr-1/parkin knockout *C. elegans*. *Neurochem. Res.* 38, 1543–1552.
- Mason, L.H., Harp, J.P., Han, D.Y., 2014. Pb neurotoxicity: neuropsychological effects of lead toxicity. *Biomed. Res. Int.* 2014, 840547.
- Masoud, A., Kiran, R., Sandhir, R., 2009. Impaired mitochondrial functions in organophosphate induced delayed neuropathy in rats. *Cell. Mol. Neurobiol.* 29, 1245–1255.
- McElwee, M.K., Freedman, J.H., 2011. Comparative toxicology of mercurials in *Caenorhabditis elegans*. *Environ. Toxicol. Chem.* 30, 2135–2141.
- Menzel, R., Yeo, H.L., Rienau, S., Li, S., Steinberg, C.E., Sturzenbaum, S.R., 2007. Cytochrome P450s and short-chain dehydrogenases mediate the toxicogenomic response of PCB52 in the nematode *Caenorhabditis elegans*. *J. Mol. Biol.* 370, 1–13.
- Miller, D.B., 1982. Neurotoxicity of the pesticidal carbamates. *Neurobehav. Toxicol. Teratol.* 4, 779–787.
- Negga, R., Rudd, D.A., Davis, N.S., Justice, A.N., Hatfield, H.E., Valente, A.L., Fields, A.S., Fitsanakis, V.A., 2011. Exposure to Mn/Zn ethylene-bis-dithiocarbamate and glyphosate pesticides leads to neurodegeneration in *Caenorhabditis elegans*. *Neurotoxicology* 32, 331–341.
- Negga, R., Stuart, J.A., Machen, M.L., Salva, J., Lizek, A.J., Richardson, S.J., Osborne, A.S., Mirallas, O., McVey, K.A., Fitsanakis, V.A., 2012. Exposure to glyphosate- and/or Mn/Zn-ethylene-bis-dithiocarbamate-containing pesticides leads to degeneration of gamma-aminobutyric acid and dopamine neurons in *Caenorhabditis elegans*. *Neurotox. Res.* 21, 281–290.
- Nguyen, J.P., Shipley, F.B., Linder, A.N., Plummer, G.S., Liu, M., Setru, S.U., Shaevitz, J.W., Leifer, A.M., 2016. Whole-brain calcium imaging with cellular resolution in freely behaving *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* 113, E1074–1081.
- Ni, M., Li, X., Yin, Z., Jiang, H., Sidoryk-Wegrzynowicz, M., Milatovic, D., Cai, J., Aschner, M., 2010. Methylmercury induces acute oxidative stress, altering Nrf2 protein level in primary microglial cells. *Toxicol. Sci.* 116, 590–603.
- O'Bryant, S.E., Edwards, M., Menon, C.V., Gong, G., Barber, R., 2011. Long-term low-level arsenic exposure is associated with poorer neuropsychological functioning: a project FRONTIER study. *Int. J. Environ. Res. Public Health* 8, 861–874.
- Olivier, K., Liu, J., Karanth, S., Zhang, H., Roane, D.S., Pope, C.N., 2001. Glucose feeding exacerbates parathion-induced neurotoxicity. *J. Toxicol. Environ. Health A* 63, 253–271.
- Olorunsogo, O.O., Bababunmi, E.A., 1980. Inhibition of succinate-linking reduction of pyridine nucleotide in rat liver mitochondria 'in vivo' by N-(phosphonomethyl)glycine. *Toxicol. Lett.* 7, 149–152.
- Olorunsogo, O.O., Bababunmi, E.A., Bassir, O., 1979. Effect of glyphosate on rat liver mitochondria in vivo. *Bull. Environ. Contam. Toxicol.* 22, 357–364.
- Ozsvath, D.L., 2009. Fluoride and environmental health: a review. *Rev. Environ. Sci. Biotechnol.* 8, 59–79.
- Peixoto, F., 2005. Comparative effects of the roundup and glyphosate on mitochondrial oxidative phosphorylation. *Chemosphere* 61, 1115–1122.
- Pentyala, S.N., Chetty, C.S., 1993. Comparative study on the changes in AChE and ATPase activities in neonate and adult rat brains under thiobencarb stress. *J. Appl. Toxicol.* 13, 39–42.
- Peres, T.V., Eying, H., Lopes, S.C., Colle, D., Goncalves, F.M., Venske, D.K., Lopes, M.W., Ben, J., Bornhorst, J., Schwerdtle, T., Aschner, M., Farina, M., Prediger, R.D., Leal, R.B., 2015. Developmental exposure to manganese induces lasting motor and

- cognitive impairment in rats. *Neurotoxicology* 50, 28–37.
- Peres, T.V., Schettinger, M.R., Chen, P., Carvalho, F., Avila, D.S., Bowman, A.B., Aschner, M., 2016. Manganese-induced neurotoxicity: a review of its behavioral consequences and neuroprotective strategies. *BMC Pharmacol. Toxicol.* 17, 57.
- Peterson, R.T., Nass, R., Boyd, W.A., Freedman, J.H., Dong, K., Narahashi, T., 2008. Use of non-mammalian alternative models for neurotoxicological study. *Neurotoxicology* 29, 546–555.
- Rahman, S.M., Kippler, M., Tofail, F., Bolte, S., Derakhshani Hamadani, J., Vahter, M., 2017. Manganese in drinking water and cognitive abilities and behavior at 10 years of age: a prospective cohort study. *Environ. Health Perspect.* 125, 057003.
- Ranpariya, V.L., Parmar, S.K., Sheth, N.R., Chandrashekar, V.M., 2011. Neuroprotective activity of *Matricaria recutita* against fluoride-induced stress in rats. *Pharm. Biol.* 49, 696–701.
- Rice, K.M., Walker Jr., E.M., Wu, M., Gillette, C., Blough, E.R., 2014. Environmental mercury and its toxic effects. *J. Prev. Med. Public Health* 47, 74–83.
- Rodriguez, M., Snoek, L.B., De Bono, M., Kammenga, J.E., 2013. Worms under stress: C. elegans stress response and its relevance to complex human disease and aging. *Trends Genet.* 29, 367–374.
- Ruan, Q.L., Ju, J.J., Li, Y.H., Li, X.B., Liu, R., Liang, G.Y., Zhang, J., Pu, Y.P., Wang, D.Y., Yin, L.H., 2012. Chlorpyrifos exposure reduces reproductive capacity owing to a damaging effect on gametogenesis in the nematode *Caenorhabditis elegans*. *J. Appl. Toxicol.* 32, 527–535.
- Rudgalvyte, M., Peltonen, J., Lakso, M., Nass, R., Wong, G., 2016. RNA-Seq reveals acute manganese exposure increases endoplasmic reticulum related and Lipocalin mRNAs in *Caenorhabditis elegans*. *J. Biochem. Mol. Toxicol.* 30, 97–105.
- Sahu, S.N., Lewis, J., Patel, I., Bozdog, S., Lee, J.H., Sprando, R., Cinar, H.N., 2013. Genomic analysis of stress response against arsenic in *Caenorhabditis elegans*. *PLoS One* 8, e66431.
- Salim, C., Rajini, P.S., 2014. Glucose feeding during development aggravates the toxicity of the organophosphorus insecticide Monocrotophos in the nematode, *Caenorhabditis elegans*. *Physiol. Behav.* 131, 142–148.
- Sanders, T., Liu, Y., Buchner, V., Tchounwou, P.B., 2009. Neurotoxic effects and biomarkers of lead exposure: a review. *Rev. Environ. Health* 24, 15–45.
- Sanfeliu, C., Sebastia, J., Cristofol, R., Rodriguez-Farre, E., 2003. Neurotoxicity of organomercurial compounds. *Neurotox. Res.* 5, 283–305.
- Sato, K., Norris, A., Sato, M., Grant, B.D., 2014. C. elegans as a model for membrane traffic. In: *WormBook: The Online Review of C. elegans Biology*, pp. 1–47.
- Savolainen, K., Hervonen, H., 1985. Dithiocarbamate fungicides decrease histochemical reactivity of cholinesterases in the gut wall of the rat. *Arch. Toxicol. Suppl.* 8, 272–276.
- Schafer, P., Muller, M., Kruger, A., Steinberg, C.E., Menzel, R., 2009. Cytochrome P450-dependent metabolism of PCB52 in the nematode *Caenorhabditis elegans*. *Arch. Biochem. Biophys.* 488, 60–68.
- Schmeisser, K., Parker, J.A., 2018 Jan. Worms on the spectrum - C. elegans models in autism research. *Exp. Neurol.* 299 (Pt A), 199–206.
- Settivari, R., Levora, J., Nass, R., 2009. The divalent metal transporter homologues SMF-1/2 mediate dopamine neuron sensitivity in *caenorhabditis elegans* models of manganism and parkinson disease. *J. Biol. Chem.* 284, 35758–35768.
- Settivari, R., VanDuyn, N., LeVora, J., Nass, R., 2013. The Nrf2/SKN-1-dependent glutathione S-transferase pi homologue GST-1 inhibits dopamine neuron degeneration in a *Caenorhabditis elegans* model of manganism. *Neurotoxicology* 38, 51–60.
- Shankar, S., Shanker, U., Shikha, 2014. Arsenic contamination of groundwater: a review of sources, prevalence, health risks, and strategies for mitigation. *Sci. World J.* 2014, 304524.
- Sharma, C., Suhalka, P., Sukhwai, P., Jaiswal, N., Bhatnagar, M., 2014. Curcumin attenuates neurotoxicity induced by fluoride: an in vivo evidence. *Pharmacogn. Mag.* 10, 61–65.
- Sidoryk-Wegrzynowicz, M., Aschner, M., 2013. Manganese toxicity in the central nervous system: the glutamine/glutamate-gamma-aminobutyric acid cycle. *J. Intern. Med.* 273, 466–477.
- Soleo, L., Defazio, G., Scarselli, R., Zefferino, R., Livrea, P., Foa, V., 1996. Toxicity of fungicides containing ethylene-bis-dithiocarbamate in serumless dissociated mesencephalic-striatal primary coculture. *Arch. Toxicol.* 70, 678–682.
- Staub, R.E., Sparks, S.E., Quistad, G.B., Casida, J.E., 1995. S-methylation as a bioactivation mechanism for mono- and dithiocarbamate pesticides as aldehyde dehydrogenase inhibitors. *Chem. Res. Toxicol.* 8, 1063–1069.
- Sun, L., Wu, Q., Liao, K., Yu, P., Cui, Q., Rui, Q., Wang, D., 2016. Contribution of heavy metals to toxicity of coal combustion related fine particulate matter (PM2.5) in *Caenorhabditis elegans* with wild-type or susceptible genetic background. *Chemosphere* 144, 2392–2400.
- Szewczyk, B., 2013. Zinc homeostasis and neurodegenerative disorders. *Front. Aging Neurosci.* 5, 33.
- Tejeda-Benitez, L., Olivero-Verbel, J., 2016. *Caenorhabditis elegans*, a biological model for research in toxicology. *Rev. Environ. Contam. Toxicol.* 237, 1–35.
- Tonkin, E.G., Valentine, H.L., Milatovic, D.M., Valentine, W.M., 2004. N,N-diethyldithiocarbamate produces copper accumulation, lipid peroxidation, and myelin injury in rat peripheral nerve. *Toxicol. Sci.* 81, 160–171.
- Tsalik, E.L., Hobert, O., 2003. Functional mapping of neurons that control locomotory behavior in *Caenorhabditis elegans*. *J. Neurobiol.* 56, 178–197.
- Tsuji, J.S., Garry, M.R., Perez, V., Chang, E.T., 2015. Low-level arsenic exposure and developmental neurotoxicity in children: a systematic review and risk assessment. *Toxicology* 337, 91–107.
- Tuschl, K., Clayton, P.T., Gospe Jr., S.M., Gulab, S., Ibrahim, S., Singhi, P., Aulakh, R., Ribeiro, R.T., Barsottini, O.G., Zaki, M.S., Del Rosario, M.L., Dyack, S., Price, V., Rideout, A., Gordon, K., Wevers, R.A., Chong, W.K., Mills, P.B., 2012. Syndrome of hepatic cirrhosis, dystonia, polycythemia, and hypermanganesemia caused by mutations in SLC30A10, a manganese transporter in man. *Am. J. Hum. Genet.* 90, 457–466.
- Mancozeb (Dithane M-45, Manzate 200)—Chemical Profile 4/87. In: University, C., University, C (Eds.), *Pesticide Management Education Program (PMEP) at Cornell University Cooperative Extension*.
- Vanduy, N., Settivari, R., Wong, G., Nass, R., 2010. SKN-1/Nrf2 inhibits dopamine neuron degeneration in a *Caenorhabditis elegans* model of methylmercury toxicity. *Toxicol. Sci.* 118, 613–624.
- Velmurugan, G., Venkatesh Babu, D.D., Ramasamy, S., 2013. Prolonged monocrotophos intake induces cardiac oxidative stress and myocardial damage in rats. *Toxicology* 307, 103–108.
- Viquez, O.M., Caito, S.W., McDonald, W.H., Friedman, D.B., Valentine, W.M., 2012. Electrophilic adduction of ubiquitin activating enzyme E1 by N,N-diethyldithiocarbamate inhibits ubiquitin activation and is accompanied by striatal injury in the rat. *Chem. Res. Toxicol.* 25, 2310–2321.
- Wani, W.Y., Gudup, S., Sunkaria, A., Bal, A., Singh, P.P., Kandimalla, R.J., Sharma, D.R., Gill, K.D., 2011. Protective efficacy of mitochondrial targeted antioxidant MitoQ against dichlorvos induced oxidative stress and cell death in rat brain. *Neuropharmacology* 61, 1193–1201.
- White, J.G., Southgate, E., Thomson, J.N., Brenner, S., 1986. The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 314, 1–340.
- Whitford, G.M., 1983. Fluorides: metabolism, mechanisms of action and safety. *Dental Hygiene* 57 (16-18, 20-12, 24-19).
- Williams, P.L., Dusenbery, D.B., 1990. A promising indicator of neurobehavioral toxicity using the nematode *Caenorhabditis elegans* and computer tracking. *Toxicol. Ind. Health* 6, 425–440.
- Woodburn, A., 2000. Glyphosate: production, pricing and use worldwide. *Pest Manag. Sci.* 56, 309–312.
- Xing, X.J., Rui, Q., Du, M., Wang, D.Y., 2009. Exposure to lead and mercury in young larvae induces more severe deficits in neuronal survival and synaptic function than in adult nematodes. *Arch. Environ. Contam. Toxicol.* 56, 732–741.
- Ye, B., Rui, Q., Wu, Q., Wang, D., 2010. Metallothioneins are required for formation of cross-adaptation response to neurobehavioral toxicity from lead and mercury exposure in nematodes. *PLoS One* 5, e14052.
- Yu, C.W., Liao, V.H., 2014. Arsenite induces neurotoxic effects on AFD neurons via oxidative stress in *Caenorhabditis elegans*. *Metallomics* 6, 1824–1831.
- Yu, C.W., How, C.M., Liao, V.H., 2016. Arsenite exposure accelerates aging process regulated by the transcription factor DAF-16/FOXO in *Caenorhabditis elegans*. *Chemosphere* 150, 632–638.
- Zhang, Y., Ye, B., Wang, D., 2010. Effects of metal exposure on associative learning behavior in nematode *Caenorhabditis elegans*. *Arch. Environ. Contam. Toxicol.* 59, 129–136.
- Zhang, Y., You, J.Z., Zhou, Y., Zhang, P.W., Qin, Q., Zhang, Z.X., 2017. The effect of dichlorvos on control of *drosophila* and its safety evaluation under different application methods. *Environ. Sci. Pollut. Res. Int.* 24, 22940–22947.